

Anti-IFN gamma Antibody
Catalog # ABO10812**Specification**

Anti-IFN gamma Antibody - Product Information

Application	WB
Primary Accession	P01579
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Interferon gamma(IFNG) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-IFN gamma Antibody - Additional Information

Gene ID 3458

Other Names

Interferon gamma, IFN-gamma, Immune interferon, IFNG

Calculated MW

19348 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Secreted.

Tissue Specificity

Released primarily from activated T lymphocytes.

Protein Name

Interferon gamma(IFN-gamma)

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human IFN gamma(151-166aa KRKRSQMLFRGRRASQ).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the type II (or gamma) interferon family.

Anti-IFN gamma Antibody - Protein Information**Name** IFNG**Function**

Type II interferon produced by immune cells such as T-cells and NK cells that plays crucial roles in antimicrobial, antiviral, and antitumor responses by activating effector immune cells and enhancing antigen presentation (PubMed: [16914093](http://www.uniprot.org/citations/16914093), PubMed: [8666937](http://www.uniprot.org/citations/8666937)). Primarily signals through the JAK-STAT pathway after interaction with its receptor IFNGR1 to affect gene regulation (PubMed: [8349687](http://www.uniprot.org/citations/8349687)). Upon IFNG binding, IFNGR1 intracellular domain opens out to allow association of downstream signaling components JAK2, JAK1 and STAT1, leading to STAT1 activation, nuclear translocation and transcription of IFNG-regulated genes. Many of the induced genes are transcription factors such as IRF1 that are able to further drive regulation of a next wave of transcription (PubMed: [16914093](http://www.uniprot.org/citations/16914093)). Plays a role in class I antigen presentation pathway by inducing a replacement of catalytic proteasome subunits with immunoproteasome subunits (PubMed: [8666937](http://www.uniprot.org/citations/8666937)). In turn, increases the quantity, quality, and repertoire of peptides for class I MHC loading (PubMed: [8163024](http://www.uniprot.org/citations/8163024)). Increases the efficiency of peptide generation also by inducing the expression of activator PA28 that associates with the proteasome and alters its proteolytic cleavage preference (PubMed: [11112687](http://www.uniprot.org/citations/11112687)). Up-regulates as well MHC II complexes on the cell surface by promoting expression of several key molecules such as cathepsins B/CTSB, H/CTSH, and L/CTSL (PubMed: [7729559](http://www.uniprot.org/citations/7729559)). Participates in the regulation of hematopoietic stem cells during development and under homeostatic conditions by affecting their development, quiescence, and differentiation (By similarity).

Cellular Location

Secreted.

Tissue Location

Released primarily from activated T lymphocytes.


Anti-IFN gamma Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IFN gamma Antibody - Images



Anti-IFN gamma antibody, ABO10812, Western blotting All lanes: Anti IFN gamma (ABO10812) at 0.5ug/ml
Lane 1: Recombinant Human IFN gamma Protein 10ng
Lane 2: Recombinant Human IFN gamma Protein 5ng
Lane 3: Recombinant Human IFN gamma Protein 2.5ng
Lane 4: Recombinant Human IFN gamma Protein 1.25ng
Predicted bind size: 19KD
Observed bind size: 19KD

Anti-IFN gamma Antibody - Background

Interferon-gamma (IFN-gamma) is an inflammatory cytokine that has been implicated in the development of fibrosis in inflamed tissues. The production of IFN-gamma, which is under genetic control, can influence the development of fibrosis in lung allografts. IFN-gamma is also produced by natural killer (NK) cells and most prominently by CD8 cytotoxic T cells, and is vital for the control of microbial pathogens. Interferon gamma is believed to be crucial for host defence against many infections. Genetically determined variability in IFN-gamma and expression might be important for the development of tuberculosis. IFN-gamma activates human macrophage oxidative metabolism and antimicrobial activity. In addition to having antiviral activity, IFN-gamma has important immunoregulatory functions. IFN-gamma plays an important role in the control of neointima proliferation.