

Anti-Caspase-10 Antibody
Catalog # ABO10831**Specification**

Anti-Caspase-10 Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q92851
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Caspase-10(CASP10) detection. Tested with WB, IHC-P in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Caspase-10 Antibody - Additional Information

Gene ID 843

Other Names

Caspase-10, CASP-10, 3.4.22.63, Apoptotic protease Mch-4, FAS-associated death domain protein interleukin-1B-converting enzyme 2, FLICE2, ICE-like apoptotic protease 4, Caspase-10 subunit p23/17, Caspase-10 subunit p12, CASP10, MCH4

Calculated MW

58951 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Tissue Specificity

Detectable in most tissues. Lowest expression is seen in brain, kidney, prostate, testis and colon.

Protein Name

Caspase-10(CASP-10)

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human CASP10(220-236aa VKTFLEALPRAAVYRMN).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the peptidase C14A family.

Anti-Caspase-10 Antibody - Protein Information

Name CASP10

Synonyms MCH4

Function

Involved in the activation cascade of caspases responsible for apoptosis execution. Recruited to both Fas- and TNFR-1 receptors in a FADD dependent manner. May participate in the granzyme B apoptotic pathways. Cleaves and activates effector caspases CASP3, CASP4, CASP6, CASP7, CASP8 and CASP9. Hydrolyzes the small- molecule substrates, Tyr- Val-Ala-Asp-|-AMC and Asp-Glu-Val-Asp-|-AMC.


Tissue Location


Detectable in most tissues. Lowest expression is seen in brain, kidney, prostate, testis and colon

Anti-Caspase-10 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Caspase-10 Antibody - Images



Anti-Caspase-10 antibody, ABO10831, Western blotting
Lane 1: COLO320 Cell Lysate
Lane 2: HELA Cell Lysate
Lane 3: SW620 Cell Lysate
Lane 4: RAJI Cell Lysate



Anti-Caspase-10 antibody, ABO10831, IHC(P)
IHC(P): Human Intestinal Cancer Tissue

Anti-Caspase-10 Antibody - Background

Caspase-10 is an enzyme that, in humans, is encoded by the CASP10 gene. The Caspase 10 gene contains 11 exons and spans about 48 kb. This gene is mapped to 2q33.1. It is transcribed in the centromere-to-telomere direction. This gene encodes a protein that is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes that undergo proteolytic processing at conserved aspartic residues to produce two subunits, large and small, that dimerize to form the active enzyme. This protein cleaves and activates caspases 3 and 7, and the protein itself is processed by caspase 8. Mutations in this gene are associated with apoptosis defects seen in type II autoimmune lymphoproliferative syndrome. Three alternatively spliced transcript variants encoding different isoforms have been described for this gene.