

Anti-Desmoglein 2 Antibody
Catalog # ABO10877**Specification**

Anti-Desmoglein 2 Antibody - Product Information

Application	WB
Primary Accession	Q14126
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Desmoglein-2(DSG2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Desmoglein 2 Antibody - Additional Information

Gene ID 1829

Other Names

Desmoglein-2, Cadherin family member 5, HDGC, DSG2, CDHF5

Calculated MW

122294 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Rat, Mouse, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse

Subcellular Localization

Cell membrane; Single-pass type I membrane protein. Cell junction, desmosome.

Tissue Specificity

All of the tissues tested and carcinomas.

Protein Name

Desmoglein-2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Desmoglein 2(1104-1118aa STRVTKHSTVQHSYS), different from the related mouse and rat sequences by one amino acid.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Contains 4 cadherin domains.

Anti-Desmoglein 2 Antibody - Protein Information

Name DSG2

Synonyms CDHF5

Function

Component of intercellular desmosome junctions. Involved in the interaction of plaque proteins and intermediate filaments mediating cell-cell adhesion.

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell junction, desmosome


Tissue Location


All of the tissues tested and carcinomas.

Anti-Desmoglein 2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Desmoglein 2 Antibody - Images



Anti-Desmoglein 2 antibody, ABO10877, Western blotting All lanes: Anti Desmoglein 2 (ABO10877) at 0.5ug/ml
Lane 1: HT1080 Whole Cell Lysate at 40ug
Lane 2: HELA Whole Cell Lysate at 40ug
Lane 3: SW620 Whole Cell Lysate at 40ug
Lane 4: SCG Whole Cell Lysate at 40ug
Lane 5: COLO320 Whole Cell Lysate at 40ug
Predicted bind size: 122KD
Observed bind size: 122KD

Anti-Desmoglein 2 Antibody - Background

Desmoglein-2 is a protein that in humans is encoded by the DSG2 gene. These desmoglein gene family members are located in a cluster on chromosome 18. This second family member is expressed in colon, colon carcinoma, and other simple and stratified epithelial-derived cell lines. Mutations in DSG2 display a high degree of penetrance. Disease expression was of variable severity with LV involvement a prominent feature. The low prevalence of classical ECG changes highlights the need to expand current diagnostic criteria to take account of LV disease, childhood disease expression, and incomplete penetrance.