

Anti-SPARC Antibody
Catalog # ABO10899**Specification**

Anti-SPARC Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P09486
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for SPARC(SPARC) detection. Tested with WB, IHC-P in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-SPARC Antibody - Additional Information

Gene ID 6678

Other Names

SPARC, Basement-membrane protein 40, BM-40, Osteonectin, ON, Secreted protein acidic and rich in cysteine, SPARC, ON

Calculated MW

34632 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Secreted, extracellular space, extracellular matrix, basement membrane . In or around the basement membrane.

Protein Name

SPARC

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human SPARC(284-303aa DEWAGCFGIKQKDIDKDLVI).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the SPARC family.

Anti-SPARC Antibody - Protein Information

Name SPARC

Synonyms ON

Function

Appears to regulate cell growth through interactions with the extracellular matrix and cytokines. Binds calcium and copper, several types of collagen, albumin, thrombospondin, PDGF and cell membranes. There are two calcium binding sites; an acidic domain that binds 5 to 8 Ca(2+) with a low affinity and an EF-hand loop that binds a Ca(2+) ion with a high affinity.


Cellular Location


Secreted, extracellular space, extracellular matrix, basement membrane. Note=In or around the basement membrane

Anti-SPARC Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-SPARC Antibody - Images



Anti-SPARC antibody, ABO10899, Western blotting
Lane 1: Recombinant Human SPARC Protein 10ng
Lane 2: Recombinant Human SPARC Protein 5ng
Lane 3: Recombinant Human SPARC Protein 2.5ng



Anti-SPARC antibody, ABO10899, Western blotting
WB: HELA Cell Lysate



Anti-SPARC antibody, ABO10899, IHC(P)IHC(P): Human Intestinal Cancer Tissue

Anti-SPARC Antibody - Background

SPARC, secreted protein acidic and rich in cysteine, also known as Osteonectin is a protein that in humans is encoded by the SPARC gene. The human SPARC gene is 26.5 kb long, and contains 10 exons and 9 introns and is located on chromosome 5q31-q33. SPARC is a glycoprotein of 40 kD. SPARC is an acidic, cysteine-rich glycoprotein consisting of a single polypeptide chain that can be broken into 4 domains: 1) an Ca^{++} binding domains near the glutamic acidic-rich region at the amino terminus(domain I), 2) a cysteine- rich(domain II), 3) a hydrophilic region(domain III) and 4) an EF hand motif at the carboxy terminus region(domain IV). Osteonectin is a glycoprotein in the bone that binds sodium. It is secreted by osteoblasts during bone formation, initiating mineralization and promoting mineral crystal formation. Osteonectin also shows affinity for collagen in addition to bone mineral calcium. A correlation between osteonectin over expression and ampullary cancers and chronic pancreatitis has been found.