

**Anti-OSM/Oncostatin M Antibody**  
**Catalog # ABO10983****Specification**

---

**Anti-OSM/Oncostatin M Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">P13725</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Oncostatin-M(OSM) detection. Tested with WB, IHC-P in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-OSM/Oncostatin M Antibody - Additional Information**

**Gene ID** 5008

**Other Names**

Oncostatin-M, OSM, OSM

**Calculated MW**

28484 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Secreted.

**Protein Name**

Oncostatin-M(OSM)

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human Oncostatin M(181-198aa ASDAFQRKLEGCRFLHGY).

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Anti-OSM/Oncostatin M Antibody - Protein Information****Name** OSM**Function**

Growth regulator. Inhibits the proliferation of a number of tumor cell lines. Stimulates proliferation of AIDS-KS cells. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells. Uses both type I OSM receptor (heterodimers composed of LIFR and IL6ST) and type II OSM receptor (heterodimers composed of OSMR and IL6ST). Involved in the maturation of fetal hepatocytes, thereby promoting liver development and regeneration (By similarity).


**Cellular Location**

Secreted.

**Anti-OSM/Oncostatin M Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-OSM/Oncostatin M Antibody - Images**

Anti-Oncostatin M antibody, ABO10983, Western blotting All lanes: Anti Oncostatin M (ABO10983)

at 0.5ug/ml Lane 1: A549 Whole Cell Lysate at 40ug Lane 2: A549 Whole Cell Lysate at 40ug Lane 3:  
HELA Whole Cell Lysate at 40ug Predicted bind size: 28KD Observed bind size: 28KD



Anti-Oncostatin M antibody, ABO10983, IHC(P) IHC(P): Human Intestinal Cancer Tissue

#### **Anti-OSM/Oncostatin M Antibody - Background**

OSM (ONCOSTATIN M) is a member of a cytokine family that includes leukemia-inhibitory factor, granulocyte colony-stimulating factor, and interleukin 6. This gene encodes a growth regulator which inhibits the proliferation of a number of tumor cell lines. It regulates cytokine production, including IL-6, G-CSF and GM-CSF from endothelial cells. OSM is mapped on 22q12.2. OSM has the ability to inhibit the growth of human A375 melanoma cells but not normal human fibroblasts. Treatment with recombinant OSM leads to the inhibition of proliferation and changes in cellular morphology of a number of tumor cell lines derived from a wide variety of tissue types. OSM also has the ability to inhibit the proliferation of murine M1 myeloid leukemic cells and can induce their differentiation into macrophage-like cells, a function shared by LIF, CSF3, and IL6. The direction of gene transcription was telomeric to centromeric, with the OSM gene located upstream of the LIF gene.