

Anti-CYP11B1 Antibody
Catalog # ABO11010**Specification**

Anti-CYP11B1 Antibody - Product Information

Application	WB
Primary Accession	P15538
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Cytochrome P450 11B1, mitochondrial(CYP11B1) detection.
Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CYP11B1 Antibody - Additional Information

Gene ID 1584

Other Names

Cytochrome P450 11B1, mitochondrial, CYPXIB1, Cytochrome P-450c11, Cytochrome P450C11, Steroid 11-beta-hydroxylase, 1.14.15.4, CYP11B1, S11BH

Calculated MW

57573 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Mitochondrion membrane.

Protein Name

Cytochrome P450 11B1, mitochondrial

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human CYP11B1(436-450aa RNFYHVPFGFGMRQC).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-CYP11B1 Antibody - Protein Information

Name CYP11B1 {ECO:0000303|PubMed:18215163, ECO:0000312|HGNC:HGNC:2591}

Function

A cytochrome P450 monooxygenase involved in the biosynthesis of adrenal corticoids (PubMed:12530636, PubMed:1518866, PubMed:1775135, PubMed:18215163, PubMed:23322723). Catalyzes a variety of reactions that are essential for many species, including detoxification, defense, and the formation of endogenous chemicals like steroid hormones. Steroid 11beta, 18- and 19-hydroxylase with preferred regioselectivity at 11beta, then 18, and lastly 19 (By similarity). Catalyzes the hydroxylation of 11-deoxycortisol and 11-deoxycorticosterone (21- hydroxyprogesterone) at 11beta position, yielding cortisol or corticosterone, respectively, but cannot produce aldosterone (PubMed:12530636, PubMed:1518866, PubMed:1775135, PubMed:18215163, PubMed:23322723). Mechanistically, uses molecular oxygen inserting one oxygen atom into a substrate for hydroxylation and reducing the second into a water molecule. Two electrons are provided by NADPH via a two- protein mitochondrial transfer system comprising flavoprotein FDXR (adrenodoxin/ferredoxin reductase) and nonheme iron-sulfur protein FDX1 or FDX2 (adrenodoxin/ferredoxin) (PubMed:18215163). Due to its lack of 18-oxidation activity, it is incapable of generating aldosterone (PubMed:23322723). Could also be involved in the androgen metabolic pathway (Probable).

Cellular Location

Mitochondrion inner membrane {ECO:0000250|UniProtKB:P14137}; Peripheral membrane protein {ECO:0000250|UniProtKB:P14137}

Tissue Location

Expressed in the zona fasciculata/reticularis of the adrenal cortex.


Anti-CYP11B1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CYP11B1 Antibody - Images



Anti-CYP11B1 antibody, ABO11010, Western blotting All lanes: Anti CYP11B1 (ABO11010) at 0.5ug/ml
Lane 1: HELA Whole Cell Lysate at 40ug
Lane 2: U87 Whole Cell Lysate at 40ug
Lane 3: MM231 Whole Cell Lysate at 40ug
Lane 4: PANC Whole Cell Lysate at 40ug
Lane 5: MM453 Whole Cell Lysate at 40ug
Lane 6: HELA Whole Cell Lysate at 40ug
Lane 7: SMMC Whole Cell Lysate at 40ug
Predicted bind size: 58KD
Observed bind size: 58KD

Anti-CYP11B1 Antibody - Background

CYP11B1 (Cytochrome p450, family 11, subfamily B, polypeptide 1), also called Steroid 11-beta-hydroxylase or P450C11, is a steroid hydroxylase found in the zona fasciculata. The CYP11B1 gene functions primarily in mitochondria in the zona fasciculata of the adrenal cortex to convert 11-deoxycortisol to cortisol and 11-deoxycorticosterone to corticosterone. CYP11B1 is a member of the cytochrome P450 superfamily of enzymes. It is mapped on 8q24.3. The CYP11B1 gene contains 9 exons and spans 6.5 kb. Using RT-PCR, Kayes-Wandover and White detected CYP11B1 mRNA in human cardiac tissue samples from left and right atria, aorta, apex, intraventricular septum, and atrioventricular node, as well as whole adult and fetal heart. Ventricles did not express CYP11B1. In patients with 11-beta-hydroxylase deficiency leading to hypertension and congenital adrenal hyperplasia, Pascoe et al. identified mutations in the CYP11B1 gene.