

Anti-PAK6 Antibody
Catalog # ABO11038**Specification**

Anti-PAK6 Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9NQU5
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Serine/threonine-protein kinase PAK 6(PAK6) detection. Tested with WB, IHC-P in Human;Rat;Mouse.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-PAK6 Antibody - Additional Information

Gene ID 106821730;56924

Other Names

Serine/threonine-protein kinase PAK 6, 2.7.11.1, PAK-5, p21-activated kinase 6, PAK-6, PAK6, PAK5

Calculated MW

74869 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Rat, Human, Mouse, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse

Subcellular Localization

Cytoplasm. Nucleus. Cotranslocates into nucleus with AR in response to androgen induction.

Tissue Specificity

Selectively expressed in brain and testis, with lower levels in multiple tissues including prostate and breast. .

Protein Name

Serine/threonine-protein kinase PAK 6

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human PAK6(16-32aa QNFQHRVHTSFDPKGK), identical to the related rat and mouse sequences.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the protein kinase superfamily. STE Ser/Thr protein kinase family. STE20 subfamily.

Anti-PAK6 Antibody - Protein Information

Name PAK6

Synonyms PAK5

Function

Serine/threonine protein kinase that plays a role in the regulation of gene transcription. The kinase activity is induced by various effectors including AR or MAP2K6/MAPKK6. Phosphorylates the DNA-binding domain of androgen receptor/AR and thereby inhibits AR-mediated transcription. Also inhibits ESR1-mediated transcription. May play a role in cytoskeleton regulation by interacting with IQGAP1. May protect cells from apoptosis through phosphorylation of BAD.

Cellular Location

Cytoplasm. Nucleus. Note=Cotranslocates into nucleus with AR in response to androgen induction


Tissue Location


Selectively expressed in brain and testis, with lower levels in multiple tissues including prostate and breast

Anti-PAK6 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-PAK6 Antibody - Images



Anti-PAK6 antibody, ABO11038, Western blotting All lanes: Anti PAK6 (ABO11038) at 0.5ug/ml
Lane 1: HELA Whole Cell Lysate at 40ug
Lane 2: 293T Whole Cell Lysate at 40ug
Lane 3: RAJI Whole Cell Lysate at 40ug
Lane 4: COLO320 Whole Cell Lysate at 40ug
Predicted bind size: 75KD
Observed bind size: 75KD



Anti-PAK6 antibody, ABO11038, IHC(P) IHC(P): Rat Brain Tissue

Anti-PAK6 Antibody - Background

Serine/threonine-protein kinase PAK 6 (p21 Protein-activated Kinase 6) is an enzyme that in humans is encoded by the PAK6 gene. The PAK6 gene is mapped to chromosome 15q15. based on an alignment of the PAK6 sequence with the genomic sequence. This gene encodes a protein that shares a high degree of sequence similarity with p21-activated kinase (PAK) family members. The proteins of this family are Rac/Cdc42-associated Ste20-like Ser/Thr protein kinases, characterized by a highly conserved amino-terminal Cdc42/Rac interactive binding (CRIB) domain and a carboxyl-terminal kinase domain. PAK kinases are implicated in the regulation of a number of cellular processes, including cytoskeleton rearrangement, apoptosis and the MAP kinase signaling pathway. The protein encoded by this gene was found to interact with androgen receptor (AR), which is a steroid hormone-dependent transcription factor that is important for male sexual differentiation and development. The p21-activated protein kinase 6 gene was found to be highly expressed in testis and prostate tissues and the encoded protein was shown to cotranslocate into the nucleus with AR in response to androgen.