

**Anti-Cullin3 Antibody**  
**Catalog # ABO11242****Specification**

---

**Anti-Cullin3 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q13618</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Cullin-3(CUL3) detection. Tested with WB in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-Cullin3 Antibody - Additional Information**

**Gene ID** 8452

**Other Names**

Cullin-3, CUL-3, CUL3, KIAA0617

**Calculated MW**

88930 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Rat, Mouse<br>

**Subcellular Localization**

Nucleus. Golgi apparatus.

**Tissue Specificity**

Widely expressed.

**Protein Name**

Cullin-3

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human Cullin 3(426-444aa DVFERYYKQHLARRLLTNK), identical to the related rat and mouse sequences.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Belongs to the cullin family.

**Anti-Cullin3 Antibody - Protein Information**

**Name** CUL3 ([HGNC:2553](#))

**Synonyms** KIAA0617

**Function**

Core component of multiple cullin-RING-based BCR (BTB-CUL3- RBX1) E3 ubiquitin-protein ligase complexes which mediate the ubiquitination and subsequent proteasomal degradation of target proteins. BCR complexes and ARIH1 collaborate in tandem to mediate ubiquitination of target proteins (PubMed: [27565346](http://www.uniprot.org/citations/27565346)). As a scaffold protein may contribute to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. The E3 ubiquitin- protein ligase activity of the complex is dependent on the neddylation of the cullin subunit and is inhibited by the association of the deneddylated cullin subunit with TIP120A/CAND1. The functional specificity of the BCR complex depends on the BTB domain-containing protein as the substrate recognition component. BCR(KLHL42) is involved in ubiquitination of KATNA1. BCR(SPOP) is involved in ubiquitination of BMI1/PCGF4, BRMS1, MACROH2A1 and DAXX, GLI2 and GLI3. Can also form a cullin-RING-based BCR (BTB-CUL3-RBX1) E3 ubiquitin-protein ligase complex containing homodimeric SPOPL or the heterodimer formed by SPOP and SPOPL; these complexes have lower ubiquitin ligase activity. BCR(KLHL9-KLHL13) controls the dynamic behavior of AURKB on mitotic chromosomes and thereby coordinates faithful mitotic progression and completion of cytokinesis. BCR(KLHL12) is involved in ER-Golgi transport by regulating the size of COPII coats, thereby playing a key role in collagen export, which is required for embryonic stem (ES) cells division: BCR(KLHL12) acts by mediating monoubiquitination of SEC31 (SEC31A or SEC31B) (PubMed: [22358839](http://www.uniprot.org/citations/22358839), PubMed: [27716508](http://www.uniprot.org/citations/27716508)). BCR(KLHL3) acts as a regulator of ion transport in the distal nephron; by mediating ubiquitination of WNK4 (PubMed: [23387299](http://www.uniprot.org/citations/23387299), PubMed: [23453970](http://www.uniprot.org/citations/23453970), PubMed: [23576762](http://www.uniprot.org/citations/23576762)). The BCR(KLHL20) E3 ubiquitin ligase complex is involved in interferon response and anterograde Golgi to endosome transport: it mediates both ubiquitination leading to degradation and 'Lys-33'-linked ubiquitination (PubMed: [20389280](http://www.uniprot.org/citations/20389280), PubMed: [21670212](http://www.uniprot.org/citations/21670212), PubMed: [21840486](http://www.uniprot.org/citations/21840486), PubMed: [24768539](http://www.uniprot.org/citations/24768539)). The BCR(KLHL21) E3 ubiquitin ligase complex regulates localization of the chromosomal passenger complex (CPC) from chromosomes to the spindle midzone in anaphase and mediates the ubiquitination of AURKB (PubMed: [19995937](http://www.uniprot.org/citations/19995937)). The BCR(KLHL22) ubiquitin ligase complex mediates monoubiquitination of PLK1, leading to PLK1 dissociation from phosphoreceptor proteins and subsequent removal from kinetochores, allowing silencing of the spindle assembly checkpoint (SAC) and chromosome segregation (PubMed: [24768539](#)).

href="http://www.uniprot.org/citations/23455478" target="\_blank">23455478</a>). The BCR(KLHL22) ubiquitin ligase complex is also responsible for the amino acid-stimulated 'Lys-48' polyubiquitination and proteasomal degradation of DEPDC5. Through the degradation of DEPDC5, releases the GATOR1 complex-mediated inhibition of the TORC1 pathway (PubMed:<a href="http://www.uniprot.org/citations/29769719" target="\_blank">29769719</a>). The BCR(KLHL25) ubiquitin ligase complex is involved in translational homeostasis by mediating ubiquitination and subsequent degradation of hypophosphorylated EIF4EBP1 (4E-BP1) (PubMed:<a href="http://www.uniprot.org/citations/22578813" target="\_blank">22578813</a>). The BCR(KLHL25) ubiquitin ligase complex is also involved in lipid synthesis by mediating ubiquitination and degradation of ACLY (PubMed:<a href="http://www.uniprot.org/citations/27664236" target="\_blank">27664236</a>). The BCR(KBTBD8) complex acts by mediating monoubiquitination of NOLC1 and TCOF1, leading to remodel the translational program of differentiating cells in favor of neural crest specification (PubMed:<a href="http://www.uniprot.org/citations/26399832" target="\_blank">26399832</a>). Involved in ubiquitination of cyclin E and of cyclin D1 (in vitro) thus involved in regulation of G1/S transition. Involved in the ubiquitination of KEAP1, ENC1 and KLHL41 (PubMed:<a href="http://www.uniprot.org/citations/15983046" target="\_blank">15983046</a>). In concert with ATF2 and RBX1, promotes degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM. The BCR(KCTD17) E3 ubiquitin ligase complex mediates ubiquitination and degradation of TCHP, a down- regulator of cilium assembly, thereby inducing ciliogenesis (PubMed:<a href="http://www.uniprot.org/citations/25270598" target="\_blank">25270598</a>). The BCR(KLHL24) E3 ubiquitin ligase complex mediates ubiquitination of KRT14, controls KRT14 levels during keratinocytes differentiation, and is essential for skin integrity (PubMed:<a href="http://www.uniprot.org/citations/27798626" target="\_blank">27798626</a>). The BCR(KLHL18) E3 ubiquitin ligase complex mediates the ubiquitination of AURKA leading to its activation at the centrosome which is required for initiating mitotic entry (PubMed:<a href="http://www.uniprot.org/citations/23213400" target="\_blank">23213400</a>). The BCR(KEAP1) E3 ubiquitin ligase complex acts as a key sensor of oxidative and electrophilic stress by mediating ubiquitination and degradation of NFE2L2/NRF2, a transcription factor regulating expression of many cytoprotective genes (PubMed:<a href="http://www.uniprot.org/citations/15601839" target="\_blank">15601839</a>, PubMed:<a href="http://www.uniprot.org/citations/16006525" target="\_blank">16006525</a>). As part of the CUL3(KBTBD6/7) E3 ubiquitin ligase complex functions mediates 'Lys-48' ubiquitination and proteasomal degradation of TIAM1 (PubMed:<a href="http://www.uniprot.org/citations/25684205" target="\_blank">25684205</a>). By controlling the ubiquitination of that RAC1 guanine exchange factors (GEF), regulates RAC1 signal transduction and downstream biological processes including the organization of the cytoskeleton, cell migration and cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/25684205" target="\_blank">25684205</a>). The BCR(KBTBD4) E3 ubiquitin ligase complex targets CoREST corepressor complex components RCOR1, KDM1A/LSD1 and HDAC2 for proteasomal degradation with RCOR1 likely to be the primary target while degradation of KDM1A and HDAC2 is likely due to their association with RCOR1 (PubMed:<a href="http://www.uniprot.org/citations/33417871" target="\_blank">33417871</a>). It also targets RCOR3, MIER2 and MIER3 for proteasomal degradation as well as associated proteins ZNF217 and RREB1 with degradation being dependent on the presence of an ELM2 domain in the target proteins (PubMed:<a href="http://www.uniprot.org/citations/36997086" target="\_blank">36997086</a>). The BCR(ARMC5) complex mediates premature transcription termination of transcripts that are unfavorably configured for transcriptional elongation by mediating ubiquitination of Pol II subunit POLR2A (PubMed:<a href="http://www.uniprot.org/citations/35687106" target="\_blank">35687106</a>, PubMed:<a href="http://www.uniprot.org/citations/38225631" target="\_blank">38225631</a>, PubMed:<a href="http://www.uniprot.org/citations/39504960" target="\_blank">39504960</a>, PubMed:<a href="http://www.uniprot.org/citations/39667934" target="\_blank">39667934</a>). Required for 'Lys-63'-linked ubiquitination of large ribosomal subunit protein MRPL12 (PubMed:<a href="http://www.uniprot.org/citations/37526061" target="\_blank">37526061</a>).

### Cellular Location

Nucleus. Golgi apparatus. Cell projection, cilium, flagellum. Cytoplasm, cytoskeleton, spindle.

Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Note=Detected along the length of the sperm flagellum and in the cytoplasm of the germ cells (PubMed:28395323). Predominantly found in the nucleus in interphase cells, found at the centrosome at late G2 or prophase, starts accumulating at the spindle poles in prometaphase and stays on the spindle poles and the mitotic spindle at metaphase (PubMed:23213400)

#### **Tissue Location**


Brain, spermatozoa, and testis (at protein level). Widely expressed.

#### **Anti-Cullin3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **Anti-Cullin3 Antibody - Images**



Anti-Cullin3 antibody, ABO11242, Western blotting  
Lane 1: HELA Cell Lysate  
Lane 2: MCF-7 Cell Lysate  
Lane 3: Rat Testis Tissue Lysate  
Lane 4: Rat Brain Tissue Lysate

#### **Anti-Cullin3 Antibody - Background**

Cullin-3(Cul 3), is a protein that in humans is encoded by the CUL3 gene. Kipreos et al.(1996) identified a conserved gene family, designated cullins, with at least 5 members in nematodes, 6 in humans, and 3 in *S. cerevisiae*. Human CUL3 is an ortholog of nematode cul3. Hartz(2012) mapped the CUL3 gene to chromosome 2q36.2 based on an alignment of the CUL3 sequence with the genomic sequence(GRCh37). Sumara et al.(2007) found that KLHL9, KLHL13, and CUL3 interacted directly in a 370-kD protein complex in HeLa cell lysates. Rondou et al.(2008) showed that interaction between KLHL12 and the CUL3 ubiquitin ligase complex directed ubiquitination of dopamine receptor D4(DRD4). KLHL12 interacted directly with CUL3 and with the polymorphic

intracellular loop-3 of D4. By immunoprecipitation analysis of HeLa cells and HCT116 human colon carcinoma cells expressing epitope-tagged proteins, Kigoshi et al.(2011) showed that KLHL7 interacted with CUL3 and ROC1(RBX1).