

**Anti-CD11b Antibody**  
**Catalog # ABO11246****Specification**

---

**Anti-CD11b Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P05555</a>
Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Integrin alpha-M(ITGAM) detection. Tested with WB in Mouse.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-CD11b Antibody - Additional Information****Other Names**

Integrin alpha-M, CD11 antigen-like family member B, CR-3 alpha chain, Cell surface glycoprotein MAC-1 subunit alpha, Leukocyte adhesion receptor MO1, CD11b, Itgam

**Calculated MW**

127481 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Mouse<br>

**Subcellular Localization**

Membrane; Single-pass type I membrane protein.

**Tissue Specificity**

Predominantly expressed in monocytes and granulocytes.

**Protein Name**

Integrin alpha-M

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg Thimerosal, 0.05mg NaN<sub>3</sub>.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of mouse CD11b(1135-1153aa KRQYKDMNEAAPQDAPPQ).

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

Storage

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

#### Sequence Similarities

Belongs to the integrin alpha chain family.

### Anti-CD11b Antibody - Protein Information

**Name** Itgam

#### Function

Integrin ITGAM/ITGB2 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (By similarity). It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is also a receptor for fibrinogen, factor X and ICAM1. It recognizes P1 and P2 peptides of fibrinogen gamma chain. Regulates neutrophil migration. In association with beta subunit ITGB2/CD18, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (By similarity). May regulate phagocytosis-induced apoptosis in extravasated neutrophils (By similarity). May play a role in mast cell development (By similarity). Required with TYROBP/DAP12 in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development (PubMed: <http://www.uniprot.org/citations/18685038> target="\_blank">18685038</a>).

#### Cellular Location

Cell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P11215}. Membrane raft {ECO:0000250|UniProtKB:P11215}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:P11215}

#### Tissue Location


Predominantly expressed in monocytes and granulocytes (PubMed:3887182, PubMed:8986723). Expressed in a subset of peritoneal mast cells (PubMed:9862668). Expressed in microglia (at protein level) (PubMed:18685038).


### Anti-CD11b Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-CD11b Antibody - Images





Anti- CD11b antibody, ABO11246, Western blotting All lanes: Anti CD11b(ABO11246) at 0.5ug/ml WB: Mouse Liver Tissue Lysate at 50ug Predicted bind size: 127KD Observed bind size: 127KD

### Anti-CD11b Antibody - Background

ITGAM(Integrin Alpha-M), also called CD11B or Mo1 ALPHA SUBUNIT(MO1A), is one protein subunit that forms the heterodimeric integrin alpha-M beta-2(alphaMbeta2) molecule. A major surface antigen family on human leukocytes includes complement receptor type 3(CR3A; also called ITGAM, Mac1, or Mo1), lymphocyte function-associated(LFA) antigen type 1(ITGAL), and p150, 95(ITGAX). By in situ hybridization, Corbi et al.(1988) demonstrated that the genes encoding the alpha subunits of LFA1(ITGAL), Mac1, and p150, 95(ITGAX), all of which are involved in leukocyte adhesion, constitute a cluster on 16p13.1-p11. Callen et al.(1991) narrowed the assignment to 16p11.2. Inflammation plays an essential role in the initiation and progression of atherosclerosis. Simon et al.(2000) presented evidence that it also has a role in vascular repair after mechanical arterial injury(i.e., percutaneous transluminal coronary angioplasty, or PTCA).