

# **Anti-ALDH2 Antibody**

**Catalog # ABO11379** 

# Specification

## **Anti-ALDH2 Antibody - Product Information**

Application WB, IHC-P
Primary Accession P05091
Host Reactivity Human, Rat
Clonality Polyclonal
Format Lyophilized

**Description** 

Rabbit IgG polyclonal antibody for Aldehyde dehydrogenase, mitochondrial(ALDH2) detection. Tested with WB, IHC-P in Human;Rat.

#### Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

#### **Anti-ALDH2 Antibody - Additional Information**

Gene ID 217

#### **Other Names**

Aldehyde dehydrogenase, mitochondrial, 1.2.1.3, ALDH class 2, ALDH-E2, ALDHI, ALDH2, ALDM

# **Calculated MW**

56381 MW KDa

#### **Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1  $\mu$ g/ml, Human, Rat, By Heat<br/>br>Western blot, 0.1-0.5  $\mu$ g/ml, Rat, Human<br/>br>

## **Subcellular Localization**

Mitochondrion matrix.

#### **Protein Name**

Aldehyde dehydrogenase, mitochondrial

#### **Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.


## **Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human ALDH2(18-33aa SAAATQAVPAPNQQPE), different from the related mouse sequence by two amino acids, and from the related rat sequence by one amino acid.

## **Purification**

Immunogen affinity purified.







**Cross Reactivity** No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Sequence Similarities** 

Belongs to the aldehyde dehydrogenase family.

## **Anti-ALDH2 Antibody - Protein Information**

Name ALDH2

Synonyms ALDM

#### **Function**

Required for clearance of cellular formaldehyde, a cytotoxic and carcinogenic metabolite that induces DNA damage.

**Cellular Location** 

Mitochondrion matrix.


# **Anti-ALDH2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **Anti-ALDH2 Antibody - Images**





Anti-ALDH2 antibody, ABO11379, Western blottingLane 1: Rat Liver Tissue LysateLane 2: Rat Intestine Tissue LysateLane 3: Rat Lung Tissue LysateLane 4: Rat Kidney Tissue Lysate



Anti-ALDH2 antibody, ABO11379, IHC(P)IHC(P): Human Liver Cancer Tissue

## Anti-ALDH2 Antibody - Background

ALDH2(Aldehyde Dehydrogenase 2 Family) is a human gene. The enzyme encoded by this gene belongs to the aldehyde dehydrogenase family of enzymes that catalyze the chemical transformation from acetaldehyde to acetic acid. Aldehyde dehydrogenase is the second enzyme of the major oxidative pathway of alcohol metabolism. Hsu et al.(1985) assigned the ALDH2 locus to chromosome 12 by means of a cDNA probe and Southern blot analysis of somatic cell hybrids. Using an unbiased proteomic search, Chen et al.(2008) identified mitochondrial ALDH2 as an enzyme whose activation correlated with reduced ischemic heart damage in rodent models. A high-throughput screen identified a small molecule activator of ALDH2, which they called Alda-1, that, when administered to rats before an ischemic event, reduced infarct size by 60%, most likely through its inhibitory effect on the formation of cytotoxic aldehydes.