

# Anti-SLC22A1 Antibody

Catalog # ABO11483

#### Specification

## Anti-SLC22A1 Antibody - Product Information

ApplicationWBPrimary Accession015245HostRabbitReactivityHumanClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Solute carrier family 22 member 1(SLC22A1) detection. Testedwith WB in Human.Human

**Reconstitution** Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

## Anti-SLC22A1 Antibody - Additional Information

Gene ID 6580

**Other Names** Solute carrier family 22 member 1, Organic cation transporter 1, hOCT1, SLC22A1, OCT1

Calculated MW 61154 MW KDa

**Application Details** Western blot, 0.1-0.5 μg/ml, Human<br>

Subcellular Localization Basolateral cell membrane ; Multi-pass membrane protein .

**Tissue Specificity** Widely expressed with high level in liver. Isoform 1 and isoform 2 are expressed in liver. Isoform 1, isoform 2, isoform 3 and isoform 4 are expressed in glial cell lines.

**Protein Name** Solute carrier family 22 member 1

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human SLC22A1(533-550aa RKAKPKENTIYLKVQTSE), different from the related mouse and rat sequences by six amino acids.

Purification



Immunogen affinity purified.

**Cross Reactivity** No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

### Anti-SLC22A1 Antibody - Protein Information

Name SLC22A1 (HGNC:10963)

Synonyms OCT1

#### Function

Electrogenic voltage-dependent transporter that mediates the transport of a variety of organic cations such as endogenous bioactive amines, cationic drugs and xenobiotics (PubMed:<a href="http://www.uniprot.org/citations/11388889" target="\_blank">11388889</a>, PubMed:<a href="http://www.uniprot.org/citations/11408531" target="\_blank">11408531</a>, PubMed:<a href="http://www.uniprot.org/citations/12439218" target=" blank">12439218</a>, PubMed:<a href="http://www.uniprot.org/citations/12719534" target=" blank">12719534</a>, PubMed:<a href="http://www.uniprot.org/citations/15389554" target=" blank">15389554</a>, PubMed:<a href="http://www.uniprot.org/citations/16263091" target=" blank">16263091</a>, PubMed:<a href="http://www.uniprot.org/citations/16272756" target="\_blank">16272756</a>, PubMed:<a href="http://www.uniprot.org/citations/16581093" target="\_blank">16581093</a>, PubMed:<a href="http://www.uniprot.org/citations/19536068" target="\_blank">19536068</a>, PubMed:<a href="http://www.uniprot.org/citations/21128598" target=" blank">21128598</a>, PubMed:<a href="http://www.uniprot.org/citations/23680637" target=" blank">23680637</a>, PubMed:<a href="http://www.uniprot.org/citations/24961373" target=" blank">24961373</a>, PubMed:<a href="http://www.uniprot.org/citations/34040533" target=" blank">34040533</a>, PubMed:<a href="http://www.uniprot.org/citations/9187257" target="\_blank">9187257</a>, PubMed:<a href="http://www.uniprot.org/citations/9260930" target="blank">9260930</a>, PubMed:<a href="http://www.uniprot.org/citations/9655880" target="\_blank">9655880</a>). Functions as a pH- and Na(+)-independent, bidirectional transporter (By similarity). Cation cellular uptake or release is driven by the electrochemical potential (i.e. membrane potential and concentration gradient) and substrate selectivity (By similarity). Hydrophobicity is a major requirement for recognition in polyvalent substrates and inhibitors (By similarity). Primarily expressed at the basolateral membrane of hepatocytes and proximal tubules and involved in the uptake and disposition of cationic compounds by hepatic and renal clearance from the blood flow (By similarity). Most likely functions as an uptake carrier in enterocytes contributing to the intestinal elimination of organic cations from the systemic circulation (PubMed: <a href="http://www.uniprot.org/citations/16263091" target=" blank">16263091</a>). Transports endogenous monoamines such as N-1-methylnicotinamide (NMN), guanidine, histamine, neurotransmitters dopamine, serotonin and adrenaline (PubMed: <a href="http://www.uniprot.org/citations/12439218" target=" blank">12439218</a>, PubMed:<a href="http://www.uniprot.org/citations/24961373" target=" blank">24961373</a>, PubMed:<a href="http://www.uniprot.org/citations/35469921" target="\_blank">35469921</a>, PubMed:<a href="http://www.uniprot.org/citations/9260930" target=" blank">9260930</a>). Also transports natural polyamines such as spermidine, agmatine and putrescine at low affinity, but relatively high turnover (PubMed:<a href="http://www.uniprot.org/citations/21128598" target=" blank">21128598</a>). Involved in the hepatic uptake of vitamin B1/thiamine, hence regulating hepatic lipid and energy metabolism (PubMed: <a href="http://www.uniprot.org/citations/24961373" target=" blank">24961373</a>). Mediates the



bidirectional transport of acetylcholine (ACh) at the apical membrane of ciliated cell in airway epithelium, thereby playing a role in luminal release of ACh from bronchial epithelium (PubMed:<a href="http://www.uniprot.org/citations/15817714" target="\_blank">15817714</a>). Transports dopaminergic neuromodulators cyclo(his-pro) and salsolinol with lower efficency (PubMed:<a href="http://www.uniprot.org/citations/17460754" target="\_blank">17460754</a>). Transports dopaminergic neuromodulators cyclo(his-pro) and salsolinol with lower efficency (PubMed:<a href="http://www.uniprot.org/citations/17460754" target="\_blank">17460754</a>). Also capable of transporting non-amine endogenous compounds such as prostaglandin E2 (PGE2) and prostaglandin F2-alpha (PGF2-alpha) (PubMed:<a

href="http://www.uniprot.org/citations/11907186" target="\_blank">11907186</a>). May contribute to the transport of cationic compounds in testes across the blood- testis-barrier (Probable). Also involved in the uptake of xenobiotics tributyImethyIammonium (TBuMA), quinidine, N-methyl-quinine (NMQ), N- methyl-quinidine (NMQD) N-(4,4-azo-n-pentyl)-quinuclidine (APQ), azidoprocainamide methoiodide (AMP), N-(4,4-azo-n-pentyl)-21- deoxyajmalinium (APDA) and 4-(4-(dimethyIamino)styryI)-N- methyIpyridinium (ASP) (PubMed:<a href="http://www.uniprot.org/citations/11408531" target="\_blank">11408531</a>, PubMed:<a

href="http://www.uniprot.org/citations/11408531" target="\_blank">11408531</a>, PubMed:<a href="http://www.uniprot.org/citations/15389554" target="\_blank">15389554</a>, PubMed:<a href="http://www.uniprot.org/citations/35469921" target="\_blank">35469921</a>, PubMed:<a href="http://www.uniprot.org/citations/9260930" target="\_blank">9260930</a>).

#### **Cellular Location**

Basolateral cell membrane; Multi-pass membrane protein. Apical cell membrane; Multi-pass membrane protein. Lateral cell membrane; Multi-pass membrane protein. Basal cell membrane; Multi-pass membrane protein. Note=Localized to the sinusoidal/basolateral membrane of hepatocytes (By similarity). Mainly localized to the basolateral membrane of renal proximal tubular cells (By similarity). However, also identified at the apical side of proximal tubular cells (PubMed:19536068). Mainly expressed at the lateral membrane of enterocytes (PubMed:16263091). Also observed at the apical side of enterocytes (PubMed:16263091). Also observed at the apical side of enterocytes (PubMed:16263091). Localized to the luminal/apical membrane of Sertoli cells (PubMed:35307651) {ECO:0000250|UniProtKB:Q63089, ECO:0000269|PubMed:15817714, ECO:0000269|PubMed:16263091, ECO:0000269|PubMed:19536068, ECO:0000269|PubMed:23680637, ECO:0000269|PubMed:35307651}

#### **Tissue Location**

Widely expressed with high level in liver (PubMed:11388889, PubMed:23680637, PubMed:9187257, PubMed:9260930). In liver, expressed around the central vein (PubMed:16263091). Expressed in kidney (PubMed:9187257, PubMed:9260930). Expressed in small intestine enterocytes (PubMed:16263091, PubMed:23680637). Localized to peritubular myoid cells, Leydig cells and moderately to the basal membrane of Sertoli cells in testes (PubMed:35307651). Expressed in tracheal and bronchial ciliated epithelium in the respiratory tract (PubMed:15817714). Also expressed in skeletal muscle, stomach, spleen, heart, placentacolon, brain, granulycytes and lympohocytes (PubMed:9187257, PubMed:9260930). [Isoform 2]: Expressed in liver and in glial cell lines. [Isoform 4]: Expressed in glial cell lines. Not expressed in liver.

#### Anti-SLC22A1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety



#### • <u>Cell Culture</u> Anti-SLC22A1 Antibody - Images



# Lysate

## Anti-SLC22A1 Antibody - Background

Solute carrier family 22(organic cation transporter), member 1, also called SLC22A1 or OCT1 is a protein that in humans is encoded by the SLC22A1 gene. This gene is mapped to 6q25.3. Polyspecific organic cation transporters in the liver, kidney, intestine, and other organs are critical for elimination of many endogenous small organic cations as well as a wide array of drugs and environmental toxins. This gene transports the polyamines spermine and spermidine. It also transports pramipexole across the basolateral membrane of the proximal tubular epithelial cells. This gene regulated by various intracellular signaling pathways including inhibition by protein kinase A activation, and endogenously activation by the calmodulin complex, the calmodulin-dependent kinase II and LCK tyrosine kinase.