

Anti-EGR2 Antibody
Catalog # ABO11487**Specification**

Anti-EGR2 Antibody - Product Information

Application	WB
Primary Accession	P11161
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for E3 SUMO-protein ligase EGR2(EGR2) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-EGR2 Antibody - Additional Information

Gene ID 1959

Other Names

E3 SUMO-protein ligase EGR2, 6.3.2.-, AT591, Early growth response protein 2, EGR-2, Zinc finger protein Krox-20, EGR2, KROX20

Calculated MW

50302 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Nucleus.

Protein Name

E3 SUMO-protein ligase EGR2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Thimerosal, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human EGR2(153-168aa TMSQTQPDLDHLYSPP), different from the related rat and mouse sequences by one amino acid.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the EGR C2H2-type zinc-finger protein family.

Anti-EGR2 Antibody - Protein Information

Name EGR2

Synonyms KROX20

Function

Sequence-specific DNA-binding transcription factor (PubMed:17717711). Plays a role in hindbrain segmentation by regulating the expression of a subset of homeobox containing genes and in Schwann cell myelination by regulating the expression of genes involved in the formation and maintenance of myelin (By similarity). Binds to two EGR2- consensus sites EGR2A (5'-CTGTAGGAG-3') and EGR2B (5'-ATGTAGGTG-3') in the HOXB3 enhancer and promotes HOXB3 transcriptional activation (By similarity). Binds to specific DNA sites located in the promoter region of HOXA4, HOXB2 and ERBB2 (By similarity). Regulates hindbrain segmentation by controlling the expression of Hox genes, such as HOXA4, HOXB3 and HOXB2, and thereby specifying odd and even rhombomeres (By similarity). Promotes the expression of HOXB3 in the rhombomere r5 in the hindbrain (By similarity). Regulates myelination in the peripheral nervous system after birth, possibly by regulating the expression of myelin proteins, such as MPZ, and by promoting the differentiation of Schwann cells (By similarity). Involved in the development of the jaw opener musculature, probably by playing a role in its innervation through trigeminal motor neurons (By similarity). May play a role in adipogenesis, possibly by regulating the expression of CEBPB (By similarity).

Cellular Location


Nucleus {ECO:0000250|UniProtKB:P08152}.


Anti-EGR2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-EGR2 Antibody - Images





Anti-EGR2 antibody, ABO11487, Western blotting Lane 1: NIH3T3 Cell Lysate Lane 2: MCF-7 Cell Lysate

Anti-EGR2 Antibody - Background

Early growth response protein 2, also called EGR2 or E3 SUMO-protein ligase EGR2, is a protein that in humans is encoded by the EGR2 gene. This gene is mapped to 10q21.3. The protein encoded by this gene is a transcription factor with three tandem C2H2-type zinc fingers. Defects in this gene are associated with Charcot-Marie-Tooth disease type 1D(CMT1D), Charcot-Marie-Tooth disease type 4E(CMT4E), and with Dejerine-Sottas syndrome(DSS). E3 SUMO-protein ligase helping SUMO1 conjugation to its coregulators NAB1 and NAB2, whose sumoylation down-regulates EGR2 own transcriptional activity.