

**Anti-IRF1 Antibody**  
**Catalog # ABO11547****Specification****Anti-IRF1 Antibody - Product Information**

Application	WB, IHC, ICC
Primary Accession	<a href="#">P10914</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Interferon regulatory factor 1(IRF1) detection. Tested with WB, IHC-P, ICC in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-IRF1 Antibody - Additional Information****Gene ID** 3659**Other Names**

Interferon regulatory factor 1, IRF-1, IRF1

**Calculated MW**

36502 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br><br>Immunocytochemistry , 0.5-1 µg/ml, Human, -<br>Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Nucleus. Cytoplasm. MYD88-associated IRF1 migrates into the nucleus more efficiently than non-MYD88- associated IRF1.

**Protein Name**

Interferon regulatory factor 1

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

**Immunogen**

A synthetic peptide corresponding to a sequence in the middle region of human IRF1(169-183aa YMQDLEVEQALTPAL).

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

**At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Belongs to the IRF family.

**Anti-IRF1 Antibody - Protein Information****Name** IRF1**Function**

Transcriptional regulator which displays a remarkable functional diversity in the regulation of cellular responses (PubMed:<a href="http://www.uniprot.org/citations/15226432" target="\_blank">15226432</a>, PubMed:<a href="http://www.uniprot.org/citations/15509808" target="\_blank">15509808</a>, PubMed:<a href="http://www.uniprot.org/citations/17516545" target="\_blank">17516545</a>, PubMed:<a href="http://www.uniprot.org/citations/17942705" target="\_blank">17942705</a>, PubMed:<a href="http://www.uniprot.org/citations/18497060" target="\_blank">18497060</a>, PubMed:<a href="http://www.uniprot.org/citations/19404407" target="\_blank">19404407</a>, PubMed:<a href="http://www.uniprot.org/citations/19851330" target="\_blank">19851330</a>, PubMed:<a href="http://www.uniprot.org/citations/22367195" target="\_blank">22367195</a>, PubMed:<a href="http://www.uniprot.org/citations/32385160" target="\_blank">32385160</a>). Regulates transcription of IFN and IFN-inducible genes, host response to viral and bacterial infections, regulation of many genes expressed during hematopoiesis, inflammation, immune responses and cell proliferation and differentiation, regulation of the cell cycle and induction of growth arrest and programmed cell death following DNA damage (PubMed:<a href="http://www.uniprot.org/citations/15226432" target="\_blank">15226432</a>, PubMed:<a href="http://www.uniprot.org/citations/15509808" target="\_blank">15509808</a>, PubMed:<a href="http://www.uniprot.org/citations/17516545" target="\_blank">17516545</a>, PubMed:<a href="http://www.uniprot.org/citations/17942705" target="\_blank">17942705</a>, PubMed:<a href="http://www.uniprot.org/citations/18497060" target="\_blank">18497060</a>, PubMed:<a href="http://www.uniprot.org/citations/19404407" target="\_blank">19404407</a>, PubMed:<a href="http://www.uniprot.org/citations/19851330" target="\_blank">19851330</a>, PubMed:<a href="http://www.uniprot.org/citations/22367195" target="\_blank">22367195</a>). Stimulates both innate and acquired immune responses through the activation of specific target genes and can act as a transcriptional activator and repressor regulating target genes by binding to an interferon-stimulated response element (ISRE) in their promoters (PubMed:<a href="http://www.uniprot.org/citations/15226432" target="\_blank">15226432</a>, PubMed:<a href="http://www.uniprot.org/citations/15509808" target="\_blank">15509808</a>, PubMed:<a href="http://www.uniprot.org/citations/17516545" target="\_blank">17516545</a>, PubMed:<a href="http://www.uniprot.org/citations/17942705" target="\_blank">17942705</a>, PubMed:<a href="http://www.uniprot.org/citations/18497060" target="\_blank">18497060</a>, PubMed:<a href="http://www.uniprot.org/citations/19404407" target="\_blank">19404407</a>, PubMed:<a href="http://www.uniprot.org/citations/19851330" target="\_blank">19851330</a>, PubMed:<a href="http://www.uniprot.org/citations/21389130" target="\_blank">21389130</a>, PubMed:<a href="http://www.uniprot.org/citations/22367195" target="\_blank">22367195</a>). Competes with the transcriptional repressor ZBED2 for binding to a common consensus sequence in gene promoters (PubMed:<a href="http://www.uniprot.org/citations/32385160" target="\_blank">32385160</a>). Its target genes for transcriptional activation activity include: genes involved in anti-viral response, such as IFN-alpha/beta, RIGI, TNFSF10/TRAIL, ZBP1, OAS1/2, PIAS1/GBP, EIF2AK2/PKR and RSAD2/viperin;

antibacterial response, such as GBP2, GBP5 and NOS2/INOS; anti-proliferative response, such as p53/TP53, LOX and CDKN1A; apoptosis, such as BBC3/PUMA, CASP1, CASP7 and CASP8; immune response, such as IL7, IL12A/B and IL15, PTGS2/COX2 and CYBB; DNA damage responses and DNA repair, such as POLQ/POLH; MHC class I expression, such as TAP1, PSMB9/LMP2, PSME1/PA28A, PSME2/PA28B and B2M and MHC class II expression, such as CIITA; metabolic enzymes, such as ACOD1/IRG1 (PubMed:<a href="http://www.uniprot.org/citations/15226432" target="\_blank">15226432</a>, PubMed:<a href="http://www.uniprot.org/citations/15509808" target="\_blank">15509808</a>, PubMed:<a href="http://www.uniprot.org/citations/17516545" target="\_blank">17516545</a>, PubMed:<a href="http://www.uniprot.org/citations/17942705" target="\_blank">17942705</a>, PubMed:<a href="http://www.uniprot.org/citations/18497060" target="\_blank">18497060</a>, PubMed:<a href="http://www.uniprot.org/citations/19404407" target="\_blank">19404407</a>, PubMed:<a href="http://www.uniprot.org/citations/19851330" target="\_blank">19851330</a>, PubMed:<a href="http://www.uniprot.org/citations/22367195" target="\_blank">22367195</a>). Represses genes involved in anti-proliferative response, such as BIRC5/survivin, CCNB1, CCNE1, CDK1, CDK2 and CDK4 and in immune response, such as FOXP3, IL4, ANXA2 and TLR4 (PubMed:<a href="http://www.uniprot.org/citations/18641303" target="\_blank">18641303</a>, PubMed:<a href="http://www.uniprot.org/citations/22200613" target="\_blank">22200613</a>). Stimulates p53/TP53-dependent transcription through enhanced recruitment of EP300 leading to increased acetylation of p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/15509808" target="\_blank">15509808</a>, PubMed:<a href="http://www.uniprot.org/citations/18084608" target="\_blank">18084608</a>). Plays an important role in immune response directly affecting NK maturation and activity, macrophage production of IL12, Th1 development and maturation of CD8+ T-cells (PubMed:<a href="http://www.uniprot.org/citations/11244049" target="\_blank">11244049</a>, PubMed:<a href="http://www.uniprot.org/citations/11846971" target="\_blank">11846971</a>, PubMed:<a href="http://www.uniprot.org/citations/11846974" target="\_blank">11846974</a>, PubMed:<a href="http://www.uniprot.org/citations/16932750" target="\_blank">16932750</a>). Also implicated in the differentiation and maturation of dendritic cells and in the suppression of regulatory T (Treg) cells development (PubMed:<a href="http://www.uniprot.org/citations/11244049" target="\_blank">11244049</a>, PubMed:<a href="http://www.uniprot.org/citations/11846971" target="\_blank">11846971</a>, PubMed:<a href="http://www.uniprot.org/citations/11846974" target="\_blank">11846974</a>, PubMed:<a href="http://www.uniprot.org/citations/16932750" target="\_blank">16932750</a>). Acts as a tumor suppressor and plays a role not only in antagonism of tumor cell growth but also in stimulating an immune response against tumor cells (PubMed:<a href="http://www.uniprot.org/citations/20049431" target="\_blank">20049431</a>).

### Cellular Location


Nucleus. Cytoplasm {ECO:0000250|UniProtKB:P15314}. Note=MYD88-associated IRF1 migrates into the nucleus more efficiently than non-MYD88-associated IRF1 {ECO:0000250|UniProtKB:P15314}

### Anti-IRF1 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-IRF1 Antibody - Images



Anti-IRF1 antibody, ABO11547, Western blotting  
Lane 1: COLO320 Cell Lysate  
Lane 2: U87 Cell Lysate  
Lane 3: HELA Cell Lysate  
Lane 4: JURKAT Cell Lysate  
Lane 5: MCF-7 Cell Lysate



Anti-IRF1 antibody, ABO11547, IHC(P)  
IHC(P): Human Intestinal Cancer Tissue



Anti-IRF1 antibody, ABO11547, ICCICC: HELA Cell

#### **Anti-IRF1 Antibody - Background**

Interferon regulatory factor 1, also known as MAR is a protein that in humans is encoded by the IRF1 gene. The IRF1 gene is mapped to chromosome 5q31.1 by pulsed-field gel electrophoresis. IRF1 encodes interferon regulatory factor 1, a member of the interferon regulatory transcription factor (IRF) family. IRF1 serves as an activator of interferons alpha and beta transcription, and in mouse it has been shown to be required for double-stranded RNA induction of these genes. IRF1 also functions as a transcription activator of genes induced by interferons alpha, beta, and gamma. Further, IRF1 has been shown to play roles in regulating apoptosis and tumor-suppression.