

Anti-CSEN Antibody
Catalog # ABO11581**Specification**

Anti-CSEN Antibody - Product Information

Application	WB
Primary Accession	Q9Y2W7
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Calsenilin(KCNIP3) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CSEN Antibody - Additional Information

Gene ID 30818

Other Names

Calsenilin, A-type potassium channel modulatory protein 3, DRE-antagonist modulator, DREAM, Kv channel-interacting protein 3, KCHIP3, KCNIP3, CSEN, DREAM, KCHIP3

Calculated MW

29231 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Cytoplasm . Cell membrane ; Lipid-anchor . Endoplasmic reticulum . Golgi apparatus. Nucleus. Also membrane-bound, associated with the plasma membrane (PubMed:15485870). In the presence of PSEN2 associated with the endoplasmic reticulum and Golgi. The sumoylated form is present only in the nucleus. .

Tissue Specificity

Highly expressed in brain. Widely expressed at lower levels. Expression levels are elevated in brain cortex regions affected by Alzheimer disease. .

Protein Name

Calsenilin

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human CSEN(1-16aa

MQPAKEVTKASDGSLL).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the recoverin family.

Anti-CSEN Antibody - Protein Information

Name KCNIP3

Synonyms CSEN, DREAM, KCHIP3

Function

Calcium-dependent transcriptional repressor that binds to the DRE element of genes including PDYN and FOS. Affinity for DNA is reduced upon binding to calcium and enhanced by binding to magnesium. Seems to be involved in nociception (By similarity).

Cellular Location

Cytoplasm. Cell membrane; Lipid-anchor. Endoplasmic reticulum. Golgi apparatus. Nucleus. Note=Also membrane-bound, associated with the plasma membrane (PubMed:15485870). In the presence of PSEN2 associated with the endoplasmic reticulum and Golgi. The sumoylated form is present only in the nucleus.


Tissue Location


Highly expressed in brain. Widely expressed at lower levels. Expression levels are elevated in brain cortex regions affected by Alzheimer disease.

Anti-CSEN Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CSEN Antibody - Images



Anti- CSEN antibody, ABO11581, Western blotting All lanes: Anti CSEN (ABO11581) at 0.5ug/ml
Lane 1: HELA Whole Cell Lysate at 40ug
Lane 2: U87 Whole Cell Lysate at 40ug
Lane 3: SHG Whole Cell Lysate at 40ug
Predicted bind size: 29KD
Observed bind size: 29KD

Anti-CSEN Antibody - Background

KCNIP3, also known as Calsenilin, is a calcium-dependent transcriptional repressor that is highly expressed in central nervous system, thyroid gland, testis, thymus, and immune cells. This gene encodes a member of the family of voltage-gated potassium(Kv) channel-interacting proteins, which belong to the neuronal calcium sensor family of proteins. KCNIP3 is mapped to 2q11.1. The encoded protein functions as a calcium-regulated transcriptional repressor, and interacts with presenilins. KCNIP3-regulated genes have a critical function in Ig subclass production and proliferation in B lymphocytes. KCNIP3 has been shown to interact with PSEN1 and PSEN2.