

Anti-FGF21 Picoband Antibody

Catalog # ABO11681

Specification

Anti-FGF21 Picoband Antibody - Product Information

ApplicationWB, EPrimary AccessionQ9NSA1HostRabbitReactivityHumanClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Fibroblast growth factor 21(FGF21) detection. Tested with WB, ELISA in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-FGF21 Picoband Antibody - Additional Information

Gene ID 26291

Other Names Fibroblast growth factor 21, FGF-21, FGF21

Calculated MW 22300 MW KDa

Application Details ELISA , 0.1-0.5 μg/ml, Human, -
Western blot, 0.1-0.5 μg/ml, Human

Subcellular Localization Secreted .

Protein Name Fibroblast growth factor 21

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen E. coli-derived human FGF21 recombinant protein (Position: H29-S209). Human FGF21 shares 80.7% amino acid (aa) sequence identity with mouse FGF21.

Purification Immunogen affinity purified.

Cross Reactivity No cross reactivity with other proteins.



Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-FGF21 Picoband Antibody - Protein Information

Name FGF21

Function

Stimulates glucose uptake in differentiated adipocytes via the induction of glucose transporter SLC2A1/GLUT1 expression (but not SLC2A4/GLUT4 expression). Activity requires the presence of KLB. Regulates systemic glucose homeostasis and insulin sensitivity.

Cellular Location Secreted {ECO:0000250|UniProtKB:Q9JJN1}.

Anti-FGF21 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-FGF21 Picoband Antibody - Images

100KD -	
70KD -	
55KD –	
35KD-	
25KD -	_
15KD -	

Western blot analysis of FGF21 expression in human placenta extract (lane 1). FGF21 at 22KD was detected using rabbit anti-FGF21 Antigen Affinity purified polyclonal antibody (Catalog # ABO11681) at 0.5 ??g/mL. The blot was developed using chemiluminescence (ECL) method .

Anti-FGF21 Picoband Antibody - Background



FGF21 (Fibroblast growth factor 21) is a protein that in humans is encoded by the FGF21 gene, which is also a member of the fibroblast growth factor (FGF) family. The FGF21 gene is mapped on 19q13.33. Using RT-PCR, Fgf21 was expressed in several types of adipose tissue in mice, including subcutaneous and epididymal fat pads and brown adipose tissue. The level of Fgf21 expression in adipose tissue was comparable to that in liver. FGF21 stimulates glucose uptake in adipocytes but not in other cell types. This effect is additive to the activity of insulin. FGF21 treatment of adipocytes is associated with phosphorylation of FRS2, a protein linking FGF receptors to the Ras/MAP kinase pathway. FGF21 also protects animals from diet-induced obesity when overexpressed in transgenic mice and lowers blood glucose and triglyceride levels when administered to diabetic rodents. Changes in Fgf21 expression due to suckling or nutritional manipulations were associated with changes in circulating free fatty acid and ketone body levels. In differentiated mouse brown adipocytes in culture, Fgf21 treatment increased the expression of thermogenic genes, caused higher total and uncoupled respiration, and enhanced glucose oxidation.