

Anti-CCL7/MCP-3 Antibody

Catalog # ABO11753

Specification

Anti-CCL7/MCP-3 Antibody - Product Information

ApplicationWB, EPrimary AccessionQ03366HostRabbitReactivityMouseClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for C-C motif chemokine 7(CCL7) detection. Tested with WB, ELISA in Mouse.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CCL7/MCP-3 Antibody - Additional Information

Gene ID 20306

Other Names C-C motif chemokine 7, Intercrine/chemokine MARC, Monocyte chemoattractant protein 3, Monocyte chemotactic protein 3, MCP-3, Protein FIC, Small-inducible cytokine A7, Ccl7, Fic, Mcp3, Scya7

Calculated MW 10999 MW KDa

Application Details Western blot, 0.1-0.5 μg/ml, Mouse, -
ELISA , 0.1-0.5 μg/ml, Mouse

Subcellular Localization Secreted.

Protein Name C-C motif chemokine 7

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived mouse MCP3 recombinant protein (Position: Q24-P97). Mouse MCP3 shares 60% and 95% amino acid (aa) sequences identity with human and rat MCP3, respectively.

Purification Immunogen affinity purified.



Cross Reactivity No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-CCL7/MCP-3 Antibody - Protein Information

Name Ccl7

Synonyms Fic, Mcp3, Scya7

Function

Chemotactic factor that attracts monocytes and eosinophils, but not neutrophils. Augments monocyte anti-tumor activity (By similarity).

Cellular Location Secreted.

Anti-CCL7/MCP-3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- Anti-CCL7/MCP-3 Antibody Images





Anti-MCP3 Picoband antibody, ABO11753-1.jpgAll lanes: Anti-MCP3(ABO11753) at 0.5ug/mlWB: Recombinant Mouse MCP3 Protein 0.5ngPredicted bind size: 36KDObserved bind size: 36KD

Anti-CCL7/MCP-3 Antibody - Background

Chemokine (C-C motif) ligand 7 (CCL7) is a small cytokine known as a chemokine that was previously called monocyte-specific chemokine 3 (MCP3). It belongs to the C-C chemokine family. By fluorescence in situ hybridization, MCP3 gene is mapped to chromosome 17q11.2-q12. MCP3 was identified as a physiologic substrate of gelatinase A. Cleaved MCP3 binds to CC-chemokine receptors-1, -2, and -3, but no longer induces calcium fluxes or promotes chemotaxis, and instead acts as a general chemokine antagonist that dampens inflammation, it has been found that matrix metalloproteinases are both effectors and regulators of the inflammatory response.