

Anti-CREB Picoband Antibody

Catalog # ABO11807

Specification

Anti-CREB Picoband Antibody - Product Information

Application WB
Primary Accession P16220
Host Rabbit
Reactivity Human, Rat
Clonality Polyclonal
Format Lyophilized

Description

Rabbit IgG polyclonal antibody for Cyclic AMP-responsive element-binding protein 1(CREB1) detection. Tested with WB in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CREB Picoband Antibody - Additional Information

Gene ID 1385

Other Names

Cyclic AMP-responsive element-binding protein 1, CREB-1, cAMP-responsive element-binding protein 1, CREB1

Calculated MW 36688 MW KDa

Application Details

Western blot, 0.1-0.5 μg/ml, Human, Rat

Subcellular Localization

Nucleus.

Protein Name

Cyclic AMP-responsive element-binding protein 1

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived human CREB recombinant protein (Position: M1-T177). Human CREB shares 99% and 98% amino acid (aa) sequences identity with mouse and rat CREB, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity



No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence SimilaritiesBelongs to the bZIP family.

Anti-CREB Picoband Antibody - Protein Information

Name CREB1

Function

Phosphorylation-dependent transcription factor that stimulates transcription upon binding to the DNA cAMP response element (CRE), a sequence present in many viral and cellular promoters (By similarity). Transcription activation is enhanced by the TORC coactivators which act independently of Ser-119 phosphorylation (PubMed:14536081). Involved in different cellular processes including the synchronization of circadian rhythmicity and the differentiation of adipose cells (By similarity). Regulates the expression of apoptotic and inflammatory response factors in cardiomyocytes in response to ERFE-mediated activation of AKT signaling (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00312, ECO:0000255|PROSITE-ProRule:PRU00978, ECO:0000269|PubMed:12552083}

Anti-CREB Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-CREB Picoband Antibody - Images





Anti-CREB Picoband antibody, ABO11807-1.jpgAll lanes: Anti CREB (ABO11807) at 0.5ug/mlWB: Recombinant Human CREB Protein 0.5ngPredicted bind size: 38KDObserved bind size: 38KD



Anti-CREB Picoband antibody, ABO11807-2.jpgAll lanes: Anti CREB (ABO11807) at 0.5ug/mlLane 1: Rat Thymus Tissue Lysate at 50ugLane 2: Rat Brain Tissue Lysate at 50ugLane 3: HT1080 Whole Cell Lysate at 40ugPredicted bind size: 43KDObserved bind size: 43KD

Anti-CREB Picoband Antibody - Background

CREB1, also known as cAMP responsive element binding protein 1, is a protein that in humans is encoded by the CREB1 gene. It is mapped to 2q33.3. This protein binds the cAMP response element, a DNA nucleotide sequence present in many viral and cellular promoters. The binding of CREB1 stimulates transcription. This protein is a CREB transcription factor that is a member of the leucine zipper family of DNA-binding proteins. This protein binds as a homodimer to the cAMP-responsive element, an octameric palindrome. The protein is phosphorylated by several protein kinases, and induces transcription of genes in response to hormonal stimulation of the cAMP pathway.