

Anti-Kv2.1 Picoband Antibody
Catalog # ABO11817**Specification****Anti-Kv2.1 Picoband Antibody - Product Information**

Application	WB, IHC
Primary Accession	Q14721
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Potassium voltage-gated channel subfamily B member 1 (KCNB1) detection. Tested with WB, IHC-P, IHC-F in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Kv2.1 Picoband Antibody - Additional Information

Gene ID 3745

Other Names

Potassium voltage-gated channel subfamily B member 1 {ECO:0000312|HGNC:HGNC:6231}, Delayed rectifier potassium channel 1, DRK1, h-DRK1, Voltage-gated potassium channel subunit Kv2.1, KCNB1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=6231)

Calculated MW

95878 MW KDa

Application Details

Immunohistochemistry(Frozen Section), 0.5-1 µg/ml, Mouse, Rat,
-
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By
Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cell membrane . Perikaryon . Cell projection, axon . Cell projection, dendrite . Membrane;
Multi-pass membrane protein. Cell junction, synapse, postsynaptic cell membrane . Cell junction,
synapse . Cell junction, synapse, synaptosome . Lateral cell membrane . Cell membrane,
sarcolemma . Localizes to high-density somatodendritic clusters and non-clustered sites on the
surface of neocortical and hippocampal pyramidal neurons in a cortical actin
cytoskeleton-dependent manner (PubMed:24477962). Localizes also to high-density clusters in the
axon initial segment (AIS), at ankyrin-G-deficient sites, on the surface of neocortical and
hippocampal pyramidal neurons (PubMed:24477962). KCNB1-containing AIS clusters localize either
in close apposition to smooth endoplasmic reticulum cisternal organelles or with GABA-A
receptor-containing synapses of hippocampal and cortical pyramidal neurons, respectively
(PubMed:24477962). Localizes to high-density clusters on the cell surface of atrial and ventricular
myocytes and at the lateral plasma membrane in epithelial cells. Localizes both to the axial and

transverse tubules (T tubule) and sarcolemma in ventricular myocytes. Associated with lipid raft domains. In cortical neurons, apoptotic injuries induce de novo plasma membrane insertion in a SNARE-dependent manner causing an apoptotic potassium current surge. .

Tissue Specificity

Expressed in neocortical pyramidal cells (PubMed:24477962). Expressed in pancreatic beta cells (at protein level) (PubMed:12403834, PubMed:14988243). Expressed in brain, heart, lung, liver, colon, kidney and adrenal gland (PubMed:19074135). Expressed in the cortex, amygdala, cerebellum, pons, thalamus, hypothalamus, hippocampus and substantia nigra (PubMed:19074135). .

Protein Name

Potassium voltage-gated channel subfamily B member 1

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E.coli-derived human Kv2.1 recombinant protein (Position: V687-I858). Human Kv2.1 shares 88% amino acid (aa) sequence identity with both mouse and rat Kv2.1.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the potassium channel family. B (Shab) (TC 1.A.1.2) subfamily. Kv2.1/KCNB1 sub-subfamily.

Anti-Kv2.1 Picoband Antibody - Protein Information

Name KCNB1 ([HGNC:6231](#))

Function

Voltage-gated potassium channel that mediates transmembrane potassium transport in excitable membranes, primarily in the brain, but also in the pancreas and cardiovascular system. Contributes to the regulation of the action potential (AP) repolarization, duration and frequency of repetitive AP firing in neurons, muscle cells and endocrine cells and plays a role in homeostatic attenuation of electrical excitability throughout the brain (PubMed:23161216). Plays also a role in the regulation of exocytosis independently of its electrical function (By similarity). Forms tetrameric potassium- selective channels through which potassium ions pass in accordance with their electrochemical gradient. The channel alternates between opened and closed conformations in response to the voltage difference across the membrane. Homotetrameric channels mediate a delayed-rectifier voltage-dependent outward potassium current that display rapid activation and slow inactivation in response to membrane depolarization (PubMed:10484328, PubMed:12560340, PubMed:12560340).

[1283219](http://www.uniprot.org/citations/1283219), PubMed:19074135, PubMed:19717558, PubMed:24901643, PubMed:8081723). Can form functional homotetrameric and heterotetrameric channels that contain variable proportions of KCNB2; channel properties depend on the type of alpha subunits that are part of the channel (By similarity). Can also form functional heterotetrameric channels with other alpha subunits that are non-conducting when expressed alone, such as KCNF1, KCNG1, KCNG3, KCNG4, KCNH1, KCNH2, KCNS1, KCNS2, KCNS3 and KCNV1, creating a functionally diverse range of channel complexes (PubMed:10484328, PubMed:11852086, PubMed:12060745, PubMed:19074135, PubMed:19717558, PubMed:24901643). Heterotetrameric channel activity formed with KCNS3 show increased current amplitude with the threshold for action potential activation shifted towards more negative values in hypoxic- treated pulmonary artery smooth muscle cells (By similarity). Channel properties are also modulated by cytoplasmic ancillary beta subunits such as AMIGO1, KCNE1, KCNE2 and KCNE3, slowing activation and inactivation rate of the delayed rectifier potassium channels (By similarity). In vivo, membranes probably contain a mixture of heteromeric potassium channel complexes, making it difficult to assign currents observed in intact tissues to any particular potassium channel family member. Major contributor to the slowly inactivating delayed- rectifier voltage-gated potassium current in neurons of the central nervous system, sympathetic ganglion neurons, neuroendocrine cells, pancreatic beta cells, cardiomyocytes and smooth muscle cells. Mediates the major part of the somatodendritic delayed-rectifier potassium current in hippocampal and cortical pyramidal neurons and sympathetic superior cervical ganglion (CGC) neurons that acts to slow down periods of firing, especially during high frequency stimulation. Plays a role in the induction of long-term potentiation (LTP) of neuron excitability in the CA3 layer of the hippocampus (By similarity). Contributes to the regulation of glucose-induced action potential amplitude and duration in pancreatic beta cells, hence limiting calcium influx and insulin secretion (PubMed:23161216). Plays a role in the regulation of resting membrane potential and contraction in hypoxia-treated pulmonary artery smooth muscle cells. May contribute to the regulation of the duration of both the action potential of cardiomyocytes and the heart ventricular repolarization QT interval. Contributes to the pronounced pro-apoptotic potassium current surge during neuronal apoptotic cell death in response to oxidative injury. May confer neuroprotection in response to hypoxia/ischemic insults by suppressing pyramidal neurons hyperexcitability in hippocampal and cortical regions (By similarity). Promotes trafficking of KCNG3, KCNH1 and KCNH2 to the cell surface membrane, presumably by forming heterotetrameric channels with these subunits (PubMed:12060745). Plays a role in the calcium-dependent recruitment and release of fusion-competent vesicles from the soma of neurons, neuroendocrine and glucose-induced pancreatic beta cells by binding key components of the fusion machinery in a pore-independent manner (By similarity).

Cellular Location

Cell membrane. Perikaryon Cell projection, axon. Cell projection, dendrite. Membrane; Multi-pass membrane protein. Postsynaptic cell membrane {ECO:0000250|UniProtKB:P15387} Synapse {ECO:0000250|UniProtKB:P15387}. Synapse, synaptosome {ECO:0000250|UniProtKB:P15387}. Lateral cell membrane {ECO:0000250|UniProtKB:P15387}. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:P15387}. Note=Localizes to high-density somatodendritic clusters and non-clustered sites on the surface of neocortical and hippocampal pyramidal neurons in a cortical actin cytoskeleton-dependent manner (PubMed:24477962). Also localizes to high-density clusters in the axon initial segment (AIS), at ankyrin-G- deficient sites, on the surface of neocortical and hippocampal pyramidal neurons (PubMed:24477962). KCNB1-containing AIS clusters localize either in close apposition to smooth endoplasmic reticulum cisternal organelles or with GABA-A

receptor-containing synapses of hippocampal and cortical pyramidal neurons, respectively (PubMed:24477962). Localizes to high-density clusters on the cell surface of atrial and ventricular myocytes and at the lateral plasma membrane in epithelial cells. Localizes both to the axial and transverse tubules (T tubule) and sarcolemma in ventricular myocytes Associated with lipid raft domains. In cortical neurons, apoptotic injuries induce de novo plasma membrane insertion in a SNARE-dependent manner causing an apoptotic potassium current surge
{ECO:0000250|UniProtKB:P15387, ECO:0000250|UniProtKB:Q03717, ECO:0000269|PubMed:12060745, ECO:0000269|PubMed:19074135, ECO:0000269|PubMed:24477962, ECO:0000269|PubMed:24901643}

Tissue Location


Expressed in neocortical pyramidal cells (PubMed:24477962). Expressed in pancreatic beta cells (at protein level) (PubMed:12403834, PubMed:14988243). Expressed in brain, heart, lung, liver, colon, kidney and adrenal gland (PubMed:19074135) Expressed in the cortex, amygdala, cerebellum, pons, thalamus, hypothalamus, hippocampus and substantia nigra (PubMed:19074135)

Anti-Kv2.1 Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-Kv2.1 Picoband Antibody - Images




Anti-Kv2.1 Picoband antibody, ABO11817-1.jpg All lanes: Anti KV2.1 (ABO11817) at 0.5ug/ml WB: Recombinant Human kv2.1 Protein 0.5ng Predicted bind size: 47KD Observed bind size: 47KD




Anti-Kv2.1 Picoband antibody, ABO11817-2.jpg All lanes: Anti KV2.1 (ABO11817) at 0.5ug/ml
Lane 1: Rat Brain Tissue Lysate at 50ug
Lane 2: Mouse Brain Tissue Lysate at 50ug
Predicted bind size: 96KD
Observed bind size: 96KD




Anti-Kv2.1 Picoband antibody, ABO11817-3.JPG IHC(P): Human Lung Cancer Tissue




Anti-Kv2.1 Picoband antibody, ABO11817-4.JPG IHC(P): Rat Brain Tissue



Anti-Kv2.1 Picoband antibody, ABO11817-5.JPGIHC(P): Mouse Brain Tissue



Anti-Kv2.1 Picoband antibody, ABO11817-6.JPGIHC(F): Rat Brain Tissue



Anti-Kv2.1 Picoband antibody, ABO11817-7.JPGIHC(F): Mouse Brain Tissue

Anti-Kv2.1 Picoband Antibody - Background

KCNB1, also known as Kv2.1 or DRK1, is a protein that, in humans, is encoded by the KCNB1 gene. It is mapped to 20q13.13. KCNB1 is found in cardiomyocytes, skeletal muscles, vascular smooth muscles, placental vasculature, retina, and pancreatic beta-cells. It can mediate the voltage-dependent potassium ion permeability of excitable membranes. KCNB1 represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume.