

Anti-HNF1 Picoband Antibody
Catalog # ABO11905**Specification****Anti-HNF1 Picoband Antibody - Product Information**

Application	WB
Primary Accession	P20823
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Hepatocyte nuclear factor 1-alpha(HNF1A) detection. Tested with WB in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-HNF1 Picoband Antibody - Additional Information

Gene ID 6927

Other Names

Hepatocyte nuclear factor 1-alpha, HNF-1-alpha, HNF-1A, Liver-specific transcription factor LF-B1, LFB1, Transcription factor 1, TCF-1, HNF1A, TCF1

Calculated MW

67356 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Rat

Subcellular Localization

Nucleus .

Tissue Specificity

Liver.

Protein Name

Hepatocyte nuclear factor 1-alpha

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived human HNF1 recombinant protein (Position: A431-Q631). Human HNF1 shares 93% and 92% amino acid (aa) sequences identity with mouse and rat HNF1, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the HNF1 homeobox family.

Anti-HNF1 Picoband Antibody - Protein Information

Name HNF1A

Synonyms TCF1

Function

Transcriptional activator that regulates the tissue specific expression of multiple genes, especially in pancreatic islet cells and in liver (By similarity). Binds to the inverted palindrome 5'-GTTAATNATTAAC-3' (PubMed: [10966642](http://www.uniprot.org/citations/10966642), PubMed: [12453420](http://www.uniprot.org/citations/12453420)). Activates the transcription of CYP1A2, CYP2E1 and CYP3A11 (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:10966642, ECO:0000269|PubMed:38018242}


Tissue Location


Liver.

Anti-HNF1 Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-HNF1 Picoband Antibody - Images



Anti- HNF1 antibody, ABO11905, Western blotting All lanes: Anti HNF1 (ABO11905) at 0.5ug/ml WB: Recombinant Human HNF1 Protein 0.5ng Predicted bind size: 45KD Observed bind size: 45KD



Anti- HNF1 antibody, ABO11905, Western blotting All lanes: Anti HNF1 (ABO11905) at 0.5ug/ml Lane 1: Rat Liver Tissue Lysate at 50ug Lane 2: Rat Kidney Tissue Lysate at 50ug Lane 3: RH35 Whole Cell Lysate at 40ug Lane 4: HELA Whole Cell Lysate at 40ug Lane 5: HEPG2 Whole Cell Lysate at 40ug Predicted bind size: 67KD Observed bind size: 67KD

Anti-HNF1 Picoband Antibody - Background

HNF1 homeobox A (hepatocyte nuclear factor 1 homeobox A), also known as HNF1A or TCF1, is a human gene. It is mapped to chromosome 12q24.3. The protein encoded by this gene is a transcription factor that is highly expressed in the liver and is involved in the regulation of the expression of several liver-specific genes. HNF1A can function as a dimer. What's more, deletion of the HNF1A activation domains or interruption of the HNF1-binding sites in the LCT promoter resulted in complete loss of transcriptional activity.