

Anti-MIF Picoband Antibody
Catalog # ABO11965**Specification****Anti-MIF Picoband Antibody - Product Information**

Application	WB, IHC-P, ICC
Primary Accession	P14174
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Macrophage migration inhibitory factor(MIF) detection. Tested with WB, IHC-P, ICC in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-MIF Picoband Antibody - Additional Information**Gene ID 4282****Other Names**

Macrophage migration inhibitory factor, MIF, 5.3.2.1, Glycosylation-inhibiting factor, GIF, L-dopachrome isomerase, L-dopachrome tautomerase, 5.3.3.12, Phenylpyruvate tautomerase, MIF, GLIF, MMIF

Calculated MW

12476 MW KDa

Application Details

Immunocytochemistry , 0.5-1 µg/ml, Human, -
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Secreted. Cytoplasm. Does not have a cleavable signal sequence and is secreted via a specialized, non- classical pathway. Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M.tuberculosis antigens.

Protein Name

Macrophage migration inhibitory factor

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E.coli-derived human MIF recombinant protein (Position: P2-A115). Human MIF shares 89% and 90% amino acid (aa) sequence identity with mouse and rat MIF respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the MIF family.

Anti-MIF Picoband Antibody - Protein Information

Name MIF {ECO:0000303|PubMed:2552447, ECO:0000312|HGNC:HGNC:7097}

Function

Pro-inflammatory cytokine involved in the innate immune response to bacterial pathogens (PubMed:15908412, PubMed:17443469, PubMed:23776208). The expression of MIF at sites of inflammation suggests a role as mediator in regulating the function of macrophages in host defense (PubMed:15908412, PubMed:17443469, PubMed:23776208). Counteracts the anti-inflammatory activity of glucocorticoids (PubMed:15908412, PubMed:17443469, PubMed:23776208). Has phenylpyruvate tautomerase and dopachrome tautomerase activity (in vitro), but the physiological substrate is not known (PubMed:11439086, PubMed:17526494). It is not clear whether the tautomerase activity has any physiological relevance, and whether it is important for cytokine activity (PubMed:11439086, PubMed:17526494).

Cellular Location

Secreted. Cytoplasm. Note=Does not have a cleavable signal sequence and is secreted via a specialized, non-classical pathway Secreted by macrophages upon stimulation by bacterial lipopolysaccharide (LPS), or by M.tuberculosis antigens

Anti-MIF Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)


- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-MIF Picoband Antibody - Images

Anti- MIF Picoband antibody, ABO11965, Western blottingAll lanes: Anti MIF (ABO11965) at 0.5ug/mlLane 1: U87 Whole Cell Lysate at 40ugLane 2: JURKAT Whole Cell Lysate at 40ugLane 3: HUT Whole Cell Lysate at 40ugLane 4: A549 Whole Cell Lysate at 40ugLane 5: HEPG2 Whole Cell Lysate at 40ugPredicted bind size: 12KDObserved bind size: 12KD



Anti- MIF Picoband antibody, ABO11965, IHC(P)IHC(P): Human Tonsil Tissue



Anti- MIF Picoband antibody, ABO11965, ICCICC: JURKAT Cell

Anti-MIF Picoband Antibody - Background

Macrophage migration inhibitory factor (MIF or MMIF), also known as GIF, is a protein that in humans is encoded by the MIF gene. It is a cytokine released by T-lymphocytes, macrophages, and the pituitary gland that serves to integrate peripheral and central inflammatory responses. MIF gene has 3 exons separated by introns of only 189 and 95 bp, and covers less than 1 kb. The localization of the human gene for MIF is to chromosome 22q11.2. MIF plays a critical role in inflammatory diseases and atherogenesis. It is also involved in cell-mediated immunity and immunoregulation. MIF plays a role in the regulation of macrophage function in host defense through the suppression of anti-inflammatory effects of glucocorticoids.