

Anti-c-Myb Picoband Antibody
Catalog # ABO11979**Specification**

Anti-c-Myb Picoband Antibody - Product Information

Application	WB
Primary Accession	P10242
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Transcriptional activator Myb(MYB) detection. Tested with WB in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-c-Myb Picoband Antibody - Additional Information

Gene ID 4602

Other Names

Transcriptional activator Myb, Proto-oncogene c-Myb, MYB

Calculated MW

72341 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Nucleus .

Protein Name

Transcriptional activator Myb

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

E.coli-derived human c-Myb recombinant protein (Position: M1-E201). Human c-Myb shares 100% amino acid (aa) sequence identity with mouse c-Myb.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Contains 3 HTH myb-type DNA-binding domains.

Anti-c-Myb Picoband Antibody - Protein Information

Name MYB

Function

Transcriptional activator; DNA-binding protein that specifically recognize the sequence 5'-YAAC[GT]G-3'. Plays an important role in the control of proliferation and differentiation of hematopoietic progenitor cells.


Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00625, ECO:0000269|PubMed:19646965}

Anti-c-Myb Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-c-Myb Picoband Antibody - Images

Anti- c-Myb Picoband antibody, ABO11979, Western blottingAll lanes: Anti c-Myb (ABO11979) at 0.5ug/mlLane 1: Human Placenta Tissue Lysate at 50ugLane 2: SMMC Whole Cell Lysate at 40ugLane 3: HELA Whole Cell Lysate at 40ugPredicted bind size: 74KDObserved bind size: 74KD

Anti-c-Myb Picoband Antibody - Background

Myb proto-oncogene protein, also known as transcriptional activator Myb, is a protein that in humans is encoded by the MYB gene. It is a member of the MYB (myeloblastosis) family of transcription factors. This gene is mapped to 6q23.3. The protein contains three domains, an N-terminal DNA-binding domain, a central transcriptional activation domain and a C-terminal domain involved in transcriptional repression. This protein plays an essential role in the regulation of hematopoiesis and may play a role in tumorigenesis, including the regulation of miR-155 in B-cells. MYB is also a critical target of MIR150.