

Anti-P2X2 Picoband Antibody
Catalog # ABO11994**Specification**

Anti-P2X2 Picoband Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9UBL9
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for P2X purinoceptor 2(P2RX2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-P2X2 Picoband Antibody - Additional Information

Gene ID 22953

Other Names

P2X purinoceptor 2, P2X2, ATP receptor, Purinergic receptor, P2RX2, P2X2

Calculated MW

51754 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cell membrane ; Multi-pass membrane protein . Localizes to the apical membranes of hair cells in the organ of Corti.

Protein Name

P2X purinoceptor 2

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived human P2X2 recombinant protein (Position: D139-L471). Human P2X2 shares 84% amino acid (aa) sequence identity with both mouse and rat P2X2.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the P2X receptor family.

Anti-P2X2 Picoband Antibody - Protein Information

Name P2RX2 ([HGNC:15459](#))

Synonyms P2X2

Function

ATP-gated nonselective transmembrane cation channel permeable to potassium, sodium and calcium (PubMed: [10570044](http://www.uniprot.org/citations/10570044), PubMed: [31636190](http://www.uniprot.org/citations/31636190)). Activation by extracellular ATP induces a variety of cellular responses, such as excitatory postsynaptic responses in sensory neurons, neuromuscular junctions (NMJ) formation, hearing, perception of taste and peristalsis (By similarity). In the inner ear, regulates sound transduction and auditory neurotransmission, outer hair cell electromotility, inner ear gap junctions, and K(+) recycling (PubMed: [23345450](http://www.uniprot.org/citations/23345450)). Mediates synaptic transmission between neurons and from neurons to smooth muscle (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein {ECO:0000250|UniProtKB:P56373}. Note=Localizes to the apical membranes of hair cells in the organ of Corti.


Tissue Location


Expressed in both the central and peripheral nervous system, as well as in the pituitary gland

Anti-P2X2 Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-P2X2 Picoband Antibody - Images



Anti- P2X2 Picoband antibody, ABO11994, Western blotting All lanes: Anti P2X2 (ABO11994) at 0.5ug/ml
Lane 1: Rat Brain Tissue Lysate at 50ug
Lane 2: Mouse Brain Tissue Lysate at 50ug
Lane 3: Human Placenta Tissue Lysate at 50ug
Lane 4: HELA Whole Cell Lysate at 40ug
Lane 5: SHG Whole Cell Lysate at 40ug
Lane 6: NEURO Whole Cell Lysate at 40ug
Lane 7: 22RV1 Whole Cell Lysate at 40ug
Lane 8: U87 Whole Cell Lysate at 40ug
Predicted bind size: 52KD
Observed bind size: 60KD



Anti- P2X2 Picoband antibody, ABO11994, IHC(P) IHC(P): Human Lung Cancer Tissue

Anti-P2X2 Picoband Antibody - Background

The P2RX2 gene encodes the P2X2 receptor, which assembles as a trimer to form a ligand-gated ion channel gated by extracellular ATP. P2X2 receptors mediate a variety of cellular responses, including excitatory postsynaptic responses in sensory neurons. The product of this gene belongs to the family of purinoceptors for ATP. P2RX2 is mapped to 12q24.33. It has been found that ATP-activated P2RX2 influenced OHC electromotility, a stimulus-induced change in hair cell length that functions as an amplifier to determine hearing sensitivity and frequency selectivity. What's more, P2RX2 channels were necessary for development of the temporary threshold shift.