

Anti-NF-kB p65 Picoband Antibody
Catalog # ABO12015**Specification**

Anti-NF-kB p65 Picoband Antibody - Product Information

Application	WB
Primary Accession	Q04207
Host	Rabbit
Reactivity	Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Transcription factor p65(RELA) detection. Tested with WB in Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-NF-kB p65 Picoband Antibody - Additional Information

Gene ID 19697

Other Names

Transcription factor p65, Nuclear factor NF-kappa-B p65 subunit, Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3, Rela, Nfkb3

Calculated MW

60212 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Mouse, Rat

Subcellular Localization

Nucleus . Cytoplasm . Colocalized with DDX1 in the nucleus upon TNF-alpha induction (By similarity). Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B). Colocalizes with GFI1 in the nucleus after lipopolysaccharide (LPS) stimulation. .

Protein Name

Transcription factor p65

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived mouse NF-kB p65 recombinant protein (Position: D291-Q479). Mouse NF-kB p65 shares 77% amino acid (aa) sequence identity with human NF-kB p65.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-NF-κB p65 Picoband Antibody - Protein Information

Name Rela

Synonyms Nfkb3

Function

NF-κappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-κappa-B is a homo- or heterodimeric complex formed by the Rel-like domain- containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52. The heterodimeric RELA-NFKB1 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. The NF-κappa-B heterodimeric RELA-NFKB1 and RELA-REL complexes, for instance, function as transcriptional activators. NF-κappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-κappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-κappa-B inhibitor (I- kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-κappa-B complex which translocates to the nucleus. The inhibitory effect of I- kappa-B on NF-κappa-B through retention in the cytoplasm is exerted primarily through the interaction with RELA. RELA shows a weak DNA- binding site which could contribute directly to DNA binding in the NF- kappa-B complex. Beside its activity as a direct transcriptional activator, it is also able to modulate promoters accessibility to transcription factors and thereby indirectly regulate gene expression (PubMed:29813070). Associates with chromatin at the NF-κappa-B promoter region via association with DDX1. Essential for cytokine gene expression in T-cells (By similarity). The NF-κappa-B homodimeric RELA- RELA complex appears to be involved in invasion-mediated activation of IL-8 expression (By similarity).

Cellular Location

Nucleus. Cytoplasm Note=Nuclear, but also found in the cytoplasm in an inactive form complexed to an inhibitor (I-kappa-B) (PubMed:21131967). Colocalized with DDX1 in the nucleus upon TNF-alpha induction (By similarity) Colocalizes with GFI1 in the nucleus after lipopolysaccharide (LPS) stimulation. {ECO:0000250|UniProtKB:Q04206, ECO:0000269|PubMed:21131967}


Anti-NF-κB p65 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)

- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-NF-kB p65 Picoband Antibody - Images



Anti- NF-kB p65 Picoband antibody, ABO12015, Western blotting All lanes: Anti NF-kB p65 (ABO12015) at 0.5ug/ml
Lane 1: PC 12 Whole Cell Lysate at 40ug
Lane 2: NRK Whole Cell Lysate at 40ug
Lane 3: HEPA Whole Cell Lysate at 40ug
Lane 4: NIH Whole Cell Lysate at 40ug
Predicted bind size: 65KD
Observed bind size: 65KD

Anti-NF-kB p65 Picoband Antibody - Background

Transcription factor p65, also known as NFKB3 or NF-kB p65, is a protein encoded by the RELA gene. NFKB is an essential transcription factor complex involved in all types of cellular processes, including cellular metabolism, chemotaxis, etc, and it may play a role in inflammatory conditions of the peripheral nervous system. Phosphorylation and acetylation of NFKB3 are crucial post-translational modifications required for NFKB activation. It has also been shown to modulate immune responses, and activation of NFKB3 is positively associated with multiple types of cancer. In addition to that, NFKB3 antagonizes TNFR1-JNK proliferative signals in epidermis and plays a nonredundant role in restraining epidermal growth.