

**Anti-RBP4 Picoband Antibody**  
**Catalog # ABO12051****Specification****Anti-RBP4 Picoband Antibody - Product Information**

Application	WB, IHC, IHC-P, E
Primary Accession	<a href="#">P02753</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Retinol-binding protein 4(RBP4) detection. Tested with WB, ELISA in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-RBP4 Picoband Antibody - Additional Information****Gene ID 5950****Other Names**

Retinol-binding protein 4, Plasma retinol-binding protein, PRBP, RBP, Plasma retinol-binding protein(1-182), Plasma retinol-binding protein(1-181), Plasma retinol-binding protein(1-179), Plasma retinol-binding protein(1-176), RBP4

**Calculated MW**

23010 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat, -<br>ELISA , 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Secreted.

**Protein Name**

Retinol-binding protein 4

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immunogen**

E.coli-derived human RBP4 recombinant protein (Position: E19-201L). Human RBP4 shares 86% amino acid (aa) sequence identity with both mouse and rat RBP4.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins.

**Storage**

**At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

**Sequence Similarities**

Belongs to the calycin superfamily. Lipocalin family.

**Anti-RBP4 Picoband Antibody - Protein Information****Name** RBP4**Function**

Retinol-binding protein that mediates retinol transport in blood plasma (PubMed:<a href="http://www.uniprot.org/citations/5541771" target="\_blank">5541771</a>). Delivers retinol from the liver stores to the peripheral tissues (Probable). Transfers the bound all-trans retinol to STRA6, that then facilitates retinol transport across the cell membrane (PubMed:<a href="http://www.uniprot.org/citations/22665496" target="\_blank">22665496</a>).

**Cellular Location**

Secreted

**Tissue Location**


Detected in blood plasma and in urine (at protein level).

**Anti-RBP4 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-RBP4 Picoband Antibody - Images**



Anti-RBP4 Picoband antibody, ABO12051, Western blotting  
All lanes: Anti RBP4 (ABO12051) at 0.5ug/ml  
Lane 1: Rat Liver Tissue Lysate at 50ug  
Lane 2: Mouse Liver Tissue Lysate at 50ug  
Lane 3: HEPG2 Whole Cell Lysate at 40ug  
Lane 4: COLO320 Whole Cell Lysate at 40ug  
Predicted bind size: 23KD  
Observed bind size: 23KD

#### Anti-RBP4 Picoband Antibody - Background

RBP4 is also known as RDCCAS. This protein belongs to the lipocalin family and is the specific carrier for retinol (vitamin A alcohol) in the blood. It delivers retinol from the liver stores to the peripheral tissues. In plasma, the RBP-retinol complex interacts with transthyretin which prevents its loss by filtration through the kidney glomeruli. A deficiency of vitamin A blocks secretion of the binding protein posttranslationally and results in defective delivery and supply to the epidermal cells.