

Anti-FRZB Picoband Antibody
Catalog # ABO12063**Specification****Anti-FRZB Picoband Antibody - Product Information**

Application	WB, E
Primary Accession	Q92765
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Secreted frizzled-related protein 3(FRZB) detection. Tested with WB, ELISA in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-FRZB Picoband Antibody - Additional Information**Gene ID 2487****Other Names**

Secreted frizzled-related protein 3, sFRP-3, Frezzled, Fritz, Frizzled-related protein 1, FrzB-1, FRZB, FIZ, FRE, FRP, FRZB1, SFRP3

Calculated MW

36254 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, -
ELISA , 0.1-0.5 µg/ml, Human

Subcellular Localization

Secreted .

Tissue Specificity

Expressed primarily in the cartilaginous cores of the long bone during embryonic and fetal development and in the appendicular skeleton (6-13 weeks). At 13 weeks of gestation, transcripts were present in early chondroblasts of the tarsal bones of the foot, the carpal bones of the hands and the epiphysis of long bones. Highly expressed in placenta and heart, followed by brain, skeletal muscle, kidney and pancreas. Weakly expressed in lung and liver.

Protein Name

Secreted frizzled-related protein 3

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E.coli-derived human FRZB recombinant protein (Position: E176-N325). Human FRZB shares 90% amino acid (aa) sequence identity with mouse FRZB.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the secreted frizzled-related protein (sFRP) family.

Anti-FRZB Picoband Antibody - Protein Information**Name** FRZB**Synonyms** FIZ, FRE, FRP, FRZB1, SFRP3**Function**

Soluble frizzled-related proteins (sFRPS) function as modulators of Wnt signaling through direct interaction with Wnts. They have a role in regulating cell growth and differentiation in specific cell types. SFRP3/FRZB appears to be involved in limb skeletogenesis. Antagonist of Wnt8 signaling. Regulates chondrocyte maturation and long bone development.

Cellular Location

Secreted.

Tissue Location


Expressed primarily in the cartilaginous cores of the long bone during embryonic and fetal development and in the appendicular skeleton (6-13 weeks). At 13 weeks of gestation, transcripts were present in early chondroblasts of the tarsal bones of the foot, the carpal bones of the hands and the epiphysis of long bones. Highly expressed in placenta and heart, followed by brain, skeletal muscle, kidney and pancreas. Weakly expressed in lung and liver

Anti-FRZB Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-FRZB Picoband Antibody - Images



Anti- FRZB Picoband antibody, ABO12063, Western blottingAll lanes: Anti FRZB (ABO12063) at 0.5ug/mlLane 1: PANC Whole Cell Lysate at 40ugLane 2: U87 Whole Cell Lysate at 40ugPredicted bind size: 36KDObserved bind size: 36KD

Anti-FRZB Picoband Antibody - Background

FRZB is also known as FRE or OS1. The protein encoded by this gene is a secreted protein that is involved in the regulation of bone development. Defects in this gene are a cause of female-specific osteoarthritis (OA) susceptibility. FRZB is a Wnt-binding protein especially important in embryonic development. It is a competitor for the cell-surface G-protein receptor Frizzled. FRZB is localized in the extracellular plasma membrane. Unlike frizzled, frzb lacks the 7 transmembrane domains normally found in G-protein-coupled receptors. It is still considered a homolog of frizzled because it contains a Cysteine Rich Domain (CRD), and because of its intracellular C-terminus which is crucial for signaling.