

# Anti-SMN1/2 Picoband Antibody

Catalog # ABO12089

### Specification

## Anti-SMN1/2 Picoband Antibody - Product Information

ApplicationWB, IHC-PPrimary Accession016637HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Survival motor neuron protein(SMN1/2) detection. Tested withWB, IHC-P in Human; Mouse; Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

### Anti-SMN1/2 Picoband Antibody - Additional Information

Gene ID 6606;6607

**Other Names** Survival motor neuron protein, Component of gems 1, Gemin-1, SMN1, SMN, SMNT

Calculated MW 31849 MW KDa

**Application Details** Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μg/ml, Human, Mouse, Rat, By Heat<br>br>Western blot, 0.1-0.5 μg/ml, Human, Mouse, Rat <br>br>

### **Subcellular Localization**

Cytoplasm. Nucleus, gem. Nucleus, Cajal body. Cytoplasmic granule. Cytoplasm, myofibril, sarcomere, Z line . Colocalizes with Actn at the Z-line of skeletal muscle (By similarity). Under stress conditions colocalizes with RPP20/POP7 in punctuated cytoplasmic granules. Colocalized and redistributed with ZPR1 from the cytoplasm to nuclear gems (Gemini of coiled bodies) and Cajal bodies. .

#### **Tissue Specificity**

Expressed in a wide variety of tissues. Expressed at high levels in brain, kidney and liver, moderate levels in skeletal and cardiac muscle, and low levels in fibroblasts and lymphocytes. Also seen at high levels in spinal cord. Present in osteoclasts and mononuclear cells (at protein level).

Protein Name Survival motor neuron protein

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.



Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human SMN1/2(22-52aa RRGTGQSDDSDIWDDTALIKAYDKAVASFKH), identical to the related mouse and rat sequences.

**Purification** Immunogen affinity purified.

**Cross Reactivity** No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

**Sequence Similarities** Belongs to the SMN family.

# Anti-SMN1/2 Picoband Antibody - Protein Information

Name SMN1

Synonyms SMN, SMNT

#### Function

The SMN complex catalyzes the assembly of small nuclear ribonucleoproteins (snRNPs), the building blocks of the spliceosome, and thereby plays an important role in the splicing of cellular pre- mRNAs (PubMed: <a href="http://www.uniprot.org/citations/18984161" target=" blank">18984161</a>, PubMed:<a href="http://www.uniprot.org/citations/9845364" target=" blank">9845364</a>). Most spliceosomal snRNPs contain a common set of Sm proteins SNRPB, SNRPD1, SNRPD2, SNRPD3, SNRPE, SNRPF and SNRPG that assemble in a heptameric protein ring on the Sm site of the small nuclear RNA to form the core snRNP (Sm core) (PubMed:<a href="http://www.uniprot.org/citations/18984161" target="\_blank">18984161</a>). In the cytosol, the Sm proteins SNRPD1, SNRPD2, SNRPE, SNRPF and SNRPG are trapped in an inactive 6S pICIn-Sm complex by the chaperone CLNS1A that controls the assembly of the core snRNP (PubMed:<a href="http://www.uniprot.org/citations/18984161" target="\_blank">18984161</a>). To assemble core snRNPs, the SMN complex accepts the trapped 5Sm proteins from CLNS1A forming an intermediate (PubMed:<a href="http://www.uniprot.org/citations/18984161" target=" blank">18984161</a>). Within the SMN complex. SMN1 acts as a structural backbone and together with GEMIN2 it gathers the Sm complex subunits (PubMed:<a href="http://www.uniprot.org/citations/17178713" target=" blank">17178713</a>, PubMed:<a href="http://www.uniprot.org/citations/21816274" target="\_blank">21816274</a>, PubMed:<a href="http://www.uniprot.org/citations/22101937" target="\_blank">22101937</a>). Binding of snRNA inside 5Sm ultimately triggers eviction of the SMN complex, thereby allowing binding of SNRPD3 and SNRPB to complete assembly of the core snRNP (PubMed:<a href="http://www.uniprot.org/citations/31799625" target="\_blank">31799625</a>). Ensures the correct splicing of U12 intron- containing genes that may be important for normal motor and proprioceptive neurons development (PubMed:<a href="http://www.uniprot.org/citations/23063131" target=" blank">23063131</a>). Also required for resolving RNA-DNA hybrids created by RNA polymerase II, that form R- loop in transcription terminal regions, an important step in proper transcription termination (PubMed:<a href="http://www.uniprot.org/citations/26700805" target=" blank">26700805</a>). May also

play a role in the metabolism of small nucleolar ribonucleoprotein (snoRNPs).

**Cellular Location** 



Nucleus, gem. Nucleus, Cajal body. Cytoplasm. Cytoplasmic granule. Perikaryon. Cell projection, neuron projection. Cell projection, axon {ECO:0000250|UniProtKB:P97801}. Cytoplasm, myofibril, sarcomere, Z line {ECO:0000250|UniProtKB:P97801}. Note=Colocalizes with actin and at the Z-line of skeletal muscle (By similarity). Under stress conditions colocalizes with RPP20/POP7 in punctuated cytoplasmic granules (PubMed:14715275). Colocalized and redistributed with ZPR1 from the cytoplasm to nuclear gems (Gemini of coiled bodies) and Cajal bodies (PubMed:11283611). Colocalizes with FMR1 in cytoplasmic granules in the soma and neurite cell processes (PubMed:18093976) {ECO:0000250|UniProtKB:P97801, ECO:0000269|PubMed:11283611, ECO:0000269|PubMed:14715275, ECO:0000269|PubMed:18093976}

### **Tissue Location**

Expressed in a wide variety of tissues. Expressed at high levels in brain, kidney and liver, moderate levels in skeletal and cardiac muscle, and low levels in fibroblasts and lymphocytes. Also seen at high levels in spinal cord. Present in osteoclasts and mononuclear cells (at protein level).

# Anti-SMN1/2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-SMN1/2 Picoband Antibody - Images



Anti- SMN1/2 Picoband antibody, ABO12089, Western blottingAll lanes: Anti SMN1/2(ABO12089) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Mouse Brain Tissue Lysate at 50ugLane 3: Rat Liver Tissue Lysate at 50ugLane 4: Mouse Liver Tissue Lysate at 50ugLane 5: 293T Whole Cell Lysate at 40ugLane 6: SMMC Whole Cell Lysate at 40ugLane 7: HEPG2 Whole Cell Lysate at 40ugLane 8: HELA Whole Cell Lysate at 40ugPredicted bind size: 32KDObserved bind size: 32KD





Anti- SMN1/2 Picoband antibody, ABO12089,IHC(P)IHC(P): Mouse Brain Tissue



Anti- SMN1/2 Picoband antibody, ABO12089,IHC(P)IHC(P): Rat Brain Tissue



Anti- SMN1/2 Picoband antibody, ABO12089,IHC(P)IHC(P): Human Mammary Cancer Tissue Anti-SMN1/2 Picoband Antibody - Background

This gene is part of a 500 kb inverted duplication on chromosome 5q13. This duplicated region contains at least four genes and repetitive elements which make it prone to rearrangements and deletions. The repetitiveness and complexity of the sequence have also caused difficulty in determining the organization of this genomic region. The telomeric and centromeric copies of this gene are nearly identical and encode the same protein. However, mutations in this gene, the telomeric copy, are associated with spinal muscular atrophy; mutations in the centromeric copy do not lead to disease. The centromeric copy may be a modifier of disease caused by mutation in the telomeric copy. The critical sequence difference between the two genes is a single nucleotide in exon 7, which is thought to be an exon splice enhancer. Note that the nine exons of both the telomeric and centromeric copies are designated historically as exon 1, 2a, 2b, and 3-8. It is



thought that gene conversion events may involve the two genes, leading to varying copy numbers of each gene. The protein encoded by this gene localizes to both the cytoplasm and the nucleus. Within the nucleus, the protein localizes to subnuclear bodies called gems which are found near coiled bodies containing high concentrations of small ribonucleoproteins (snRNPs). This protein forms heteromeric complexes with proteins such as SIP1 and GEMIN4, and also interacts with several proteins known to be involved in the biogenesis of snRNPs, such as hnRNP U protein and the small nucleolar RNA binding protein. Multiple transcript variants encoding distinct isoforms have been described.