

# Anti-Heparanase 1 Picoband Antibody

Catalog # ABO12117

## Specification

# Anti-Heparanase 1 Picoband Antibody - Product Information

ApplicationWBPrimary Accession09Y251HostRabbitReactivityHuman, RatClonalityPolyclonalFormatLyophilizedDescriptionPabbit IgG polyclonal antibody for Heparapase(HPSE) detection. T

Rabbit IgG polyclonal antibody for Heparanase(HPSE) detection. Tested with WB in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

## Anti-Heparanase 1 Picoband Antibody - Additional Information

Gene ID 10855

**Other Names** Heparanase, 3.2.1.166, Endo-glucoronidase, Heparanase-1, Hpa1, Heparanase 8 kDa subunit, Heparanase 50 kDa subunit, HPSE, HEP, HPA, HPA1, HPR1, HPSE1, HSE1

Calculated MW 61149 MW KDa

**Application Details** Western blot, 0.1-0.5 μg/ml, Human, Rat<br>

#### Subcellular Localization

Lysosome membrane; Peripheral membrane protein. Secreted. Nucleus. Proheparanase is secreted via vesicles of the Golgi. Interacts with cell membrane heparan sulfate proteoglycans (HSPGs). Endocytosed and accumulates in endosomes. Transferred to lysosomes where it is proteolytically cleaved to produce the active enzyme. Under certain stimuli, transferred to the cell surface. Associates with lipid rafts. Colocalizes with SDC1 in endosomal/lysosomal vesicles. Accumulates in perinuclear lysosomal vesicles. Heparin retains proheparanase in the extracellular medium (By similarity).

#### **Tissue Specificity**

Highly expressed in placenta and spleen and weakly expressed in lymph node, thymus, peripheral blood leukocytes, bone marrow, endothelial cells, fetal liver and tumor tissues. Also expressed in hair follicles, specifically in both Henle's and Huxley's layers of inner the root sheath (IRS) at anagen phase.

Protein Name Heparanase



**Contents** Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human Heparanase 1 (301-331aa NGRTATKEDFLNPDVLDIFISSVQKVFQVVE), different from the related mouse and rat sequences by eight amino acids.

**Purification** Immunogen affinity purified.

**Cross Reactivity** No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

## Anti-Heparanase 1 Picoband Antibody - Protein Information

Name HPSE

Synonyms HEP, HPA, HPA1, HPR1, HPSE1, HSE1

#### Function

Endoglycosidase that cleaves heparan sulfate proteoglycans (HSPGs) into heparan sulfate side chains and core proteoglycans. Participates in extracellular matrix (ECM) degradation and remodeling. Selectively cleaves the linkage between a glucuronic acid unit and an N-sulfo glucosamine unit carrying either a 3-O-sulfo or a 6-O-sulfo group. Can also cleave the linkage between a glucuronic acid unit and an N-sulfo glucosamine unit carrying a 2-O-sulfo group, but not linkages between a glucuronic acid unit and a 2-O-sulfated iduronic acid moiety. It is essentially inactive at neutral pH but becomes active under acidic conditions such as during tumor invasion and in inflammatory processes. Facilitates cell migration associated with metastasis, wound healing and inflammation. Enhances shedding of syndecans, and increases endothelial invasion and angiogenesis in myelomas. Acts as a procoagulant by increasing the generation of activation factor X in the presence of tissue factor and activation factor VII. Increases cell adhesion to the extracellular matrix (ECM), independent of its enzymatic activity. Induces AKT1/PKB phosphorylation via lipid rafts increasing cell mobility and invasion. Heparin increases this AKT1/PKB activation. Regulates osteogenesis. Enhances angiogenesis through up-regulation of SRC-mediated activation of VEGF. Implicated in hair follicle inner root sheath differentiation and hair homeostasis.

#### **Cellular Location**

Lysosome membrane; Peripheral membrane protein. Secreted. Nucleus. Note=Proheparanase is secreted via vesicles of the Golgi. Interacts with cell membrane heparan sulfate proteoglycans (HSPGs). Endocytosed and accumulates in endosomes. Transferred to lysosomes where it is proteolytically cleaved to produce the active enzyme. Under certain stimuli, transferred to the cell surface Associates with lipid rafts. Colocalizes with SDC1 in endosomal/lysosomal vesicles. Accumulates in perinuclear lysosomal vesicles. Heparin retains proheparanase in the extracellular medium (By similarity).

#### **Tissue Location**

Highly expressed in placenta and spleen and weakly expressed in lymph node, thymus, peripheral blood leukocytes, bone marrow, endothelial cells, fetal liver and tumor tissues. Also expressed in



hair follicles, specifically in both Henle's and Huxley's layers of inner the root sheath (IRS) at anagen phase

# Anti-Heparanase 1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

## Anti-Heparanase 1 Picoband Antibody - Images

116KD - 1 2 3 97KD -58KD -40KD -29KD -20KD -14KD -

Anti- Heparanase 1 Picoband antibody, ABO12117, Western blottingAll lanes: Anti Heparanase 1 (ABO12117) at 0.5ug/mlLane 1: Rat Liver Tissue Lysate at 50ugLane 2: Human Placenta Tissue Lysate at 50ugLane 3: A549 Whole Cell Lysate at 40ugPredicted bind size: 61KDObserved bind size: 61KD

## Anti-Heparanase 1 Picoband Antibody - Background

Heparanase, also known as HPSE, is an enzyme that acts both at the cell-surface and within the extracellular matrix to degrade polymeric heparan sulfate molecules into shorter chain length oligosaccharides. Heparanase is an endo-beta-D-glucuronidase capable of cleaving heparan sulfate and has been implicated in inflammation and tumor angiogenesis and metastasis. The successful penetration of the endothelial cell layer that lines the interior surface of blood vessels is an important process in the formation of blood borne tumour metastases. Heparan sulfate proteoglycans are major constituents of this layer and it has been shown that increased metastatic potential corresponds with increased heparanase activity for a number of cell lines.