

Anti-SLC2A2 Picoband Antibody

Catalog # ABO12126

Specification

Anti-SLC2A2 Picoband Antibody - Product Information

ApplicationWB, IHC-PPrimary AccessionP11168HostRabbitReactivityHumanClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Solute carrier family 2, facilitated glucose transporter member2(SLC2A2) detection. Tested with WB, IHC-P in Human.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-SLC2A2 Picoband Antibody - Additional Information

Gene ID 6514

Other Names Solute carrier family 2, facilitated glucose transporter member 2, Glucose transporter type 2, liver, GLUT-2, SLC2A2, GLUT2

Calculated MW 57490 MW KDa

Application Details Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat
blot, 0.1-0.5 μ g/ml, Human
br>

Subcellular Localization Membrane; Multi-pass membrane protein.

Tissue Specificity Liver, insulin-producing beta cell, small intestine and kidney.

Protein Name Solute carrier family 2, facilitated glucose transporter member 2

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human SLC2A2 (486-514aa ETKGKSFEEIAAEFQKKSGSAHRPKAAVE), different from the related mouse sequence by five amino acids, and from the related rat sequence by seven amino acids.



Purification Immunogen affinity purified.

Cross Reactivity No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the major facilitator superfamily. Sugar transporter (TC 2.A.1.1) family. Glucose transporter subfamily.

Anti-SLC2A2 Picoband Antibody - Protein Information

Name SLC2A2 (HGNC:11006)

Function

Facilitative hexose transporter that mediates the transport of glucose, fructose and galactose (PubMed:16186102, PubMed:23396969, PubMed:28083649, PubMed:28083649, PubMed:8027028, PubMed:8027028, PubMed:8027028, PubMed:8457197). Likely mediates the bidirectional transfer of glucose across the plasma membrane of hepatocytes and is responsible for uptake of glucose by the beta cells; may comprise part of the glucose-sensing mechanism of the beta cell (PubMed:8027028). May also participate with the Na(+)/glucose cotransporter in the transcellular transport of glucose in the small intestine and kidney (PubMed:3399500). Also able to mediate the transport of dehydroascorbate (PubMed:23396969).

Cellular Location

Cell membrane; Multi-pass membrane protein

Tissue Location

Liver, insulin-producing beta cell, small intestine and kidney.

Anti-SLC2A2 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>



Anti-SLC2A2 Picoband Antibody - Images



Anti- SLC2A2 Picoband antibody, ABO12126, Western blottingAll lanes: Anti SLC2A2 (ABO12126) at 0.5ug/mlLane 1: PANC Whole Cell Lysate at 40ugLane 2: A549 Whole Cell Lysate at 40ugLane 3: HT1080 Whole Cell Lysate at 40ugPredicted bind size: 57KDObserved bind size: 50KD



Anti- SLC2A2 Picoband antibody, ABO12126, IHC(P)IHC(P): Human Liver Cancer Tissue

Anti-SLC2A2 Picoband Antibody - Background

SLC2A2, also known as Glucose transporter 2 (GLUT2), is a transmembrane carrier protein that enables protein facilitatedglucose movement across cell membranes. This gene encodes an integral plasma membrane glycoprotein of the liver, islet beta cells, intestine, and kidney epithelium. The encoded protein mediates facilitated bidirectional glucose transport. Because of its low affinity for glucose, it has been suggested as a glucose sensor. Mutations in this gene are associated with susceptibility to diseases, including Fanconi-Bickel syndrome and noninsulin-dependent diabetes mellitus (NIDDM). Alternative splicing results in multiple transcript variants of this gene.