

Anti-Vitronectin Picoband Antibody
Catalog # ABO12147**Specification**

Anti-Vitronectin Picoband Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P04004
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Vitronectin(VTN) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Vitronectin Picoband Antibody - Additional Information

Gene ID 7448

Other Names

Vitronectin, VN, S-protein, Serum-spreading factor, V75, Vitronectin V65 subunit, Vitronectin V10 subunit, Somatomedin-B, VTN

Calculated MW

54306 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Secreted, extracellular space.

Tissue Specificity

Plasma.

Protein Name

Vitronectin

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Vitronectin (446-472 aa RVNLRTRRVDTVDPPYPRSIAQYWLGC), different from the related mouse sequence by two amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Contains 4 hemopexin repeats.

Anti-Vitronectin Picoband Antibody - Protein Information**Name** VTN**Function**

Vitronectin is a cell adhesion and spreading factor found in serum and tissues. Vitronectin interact with glycosaminoglycans and proteoglycans. Is recognized by certain members of the integrin family and serves as a cell-to-substrate adhesion molecule. Inhibitor of the membrane-damaging effect of the terminal cytolytic complement pathway.

Cellular Location

Secreted, extracellular space


Tissue Location

Expressed in the retina pigment epithelium (at protein level) (PubMed:25136834). Expressed in plasma (at protein level) (PubMed:2448300). Expressed in serum (at protein level) (PubMed:29567995).

Anti-Vitronectin Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Vitronectin Picoband Antibody - Images



Anti- Vitronectin Picoband antibody, ABO12147, Western blotting All lanes: Anti Vitronectin (ABO12147) at 0.5ug/ml
Lane 1: Rat Liver Tissue Lysate at 50ug
Lane 2: Rat Lung Tissue Lysate at 50ug
Lane 3: HEPG2 Whole Cell Lysate at 40ug
Lane 4: HELA Whole Cell Lysate at 40ug
Lane 5: HEPA Whole Cell Lysate at 40ug
Predicted bind size: 54KD
Observed bind size: 54KD



Anti- Vitronectin Picoband antibody, ABO12147, IHC(P) IHC(P): Human Lung Cancer Tissue

Anti-Vitronectin Picoband Antibody - Background

Vitronectin, also known as VTN, is a protein that in humans is encoded by the VTN gene. The protein encoded by this gene is a member of the pexin family. It is found in serum and tissues and promotes cell adhesion and spreading, inhibits the membrane-damaging effect of the terminal cytolytic complement pathway, and binds to several serpin serine protease inhibitors. It is a secreted protein and exists in either a single chain form or a clipped, two chain form held together by a disulfide bond. Also vitronectin serves to regulate proteolysis initiated by plasminogen activation. In addition, vitronectin is a component of platelets and is, thus, involved in hemostasis.