

# Anti-Aquaporin 2 Picoband Antibody

Catalog # ABO12162

Specification

# Anti-Aquaporin 2 Picoband Antibody - Product Information

ApplicationWB, IHC-PPrimary AccessionP41181HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Aquaporin-2(AQP2) detection. Tested with WB, IHC-P inHuman;Mouse;Rat.Human;Mouse;Rat.

**Reconstitution** Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

# Anti-Aquaporin 2 Picoband Antibody - Additional Information

Gene ID 359

**Other Names** Aquaporin-2, AQP-2, ADH water channel, Aquaporin-CD, AQP-CD, Collecting duct water channel protein, WCH-CD, Water channel protein for renal collecting duct, AQP2

Calculated MW 28837 MW KDa

**Application Details** Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μg/ml, Human, Mouse, Rat, By Heat<br>br>Western blot, 0.1-0.5 μg/ml, Human, Mouse, Rat<br>br>

## **Subcellular Localization**

Apical cell membrane ; Multi-pass membrane protein . Basolateral cell membrane ; Multi-pass membrane protein . Cytoplasmic vesicle membrane ; Multi- pass membrane protein . Golgi apparatus, trans-Golgi network membrane ; Multi-pass membrane protein . Shuttles from vesicles to the apical membrane. Vasopressin-regulated phosphorylation is required for translocation to the apical cell membrane. PLEKHA8/FAPP2 is required to transport AQP2 from the TGN to sites where AQP2 is phosphorylated.

**Tissue Specificity** Expressed in renal collecting tubules.

Protein Name Aquaporin-2

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.



#### Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Aquaporin 2 (241-271aa EPDTDWEEREVRRRQSVELHSPQSLPRGTKA), different from the related mouse and rat sequences by one amino acid.

**Purification** Immunogen affinity purified.

**Cross Reactivity** No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Sequence Similarities Belongs to the MIP/aquaporin (TC 1.A.8) family.

## Anti-Aquaporin 2 Picoband Antibody - Protein Information

## Name AQP2 (HGNC:634)

Function

Forms a water-specific channel that provides the plasma membranes of renal collecting duct with high permeability to water, thereby permitting water to move in the direction of an osmotic gradient (PubMed:<a href="http://www.uniprot.org/citations/15509592" target="\_blank">15509592</a>, PubMed:<a href="http://www.uniprot.org/citations/7510718" target="\_blank">7510718</a>, PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">7524315</a>, PubMed:<a href="http://www.uniprot.org/citations/8140421" target="\_blank">8140421</a>, PubMed:<a href="http://www.uniprot.org/citations/8584435" target="\_blank">8584435</a>). Plays an essential role in renal water homeostasis (PubMed:<a href="http://www.uniprot.org/citations/8584435" target="\_blank">8584435</a>). Plays an essential role in renal water homeostasis (PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">8584435</a>). Plays an essential role in renal water homeostasis (PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">15509592</a>, PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">15509592</a>, PubMed:<a href="http://www.uniprot.org/citations/8584435" target="\_blank">15509592</a>, PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">15509592</a>, PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">15509592</a>, PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">8140421</a>, PubMed:<a href="http://www.uniprot.org/citations/7524315" target="\_blank">8140421</a>, PubMed:<a href="http://www.uniprot.org/citations/8584435" target="\_blank">8140421</a>). Could also be permeable to glycerol (PubMed:<a href="http://www.uniprot.org/citations/8584435" target="\_blank">8140421</a>). Could also be permeable to glycerol (PubMed:<a href="http://www.uniprot.org/citations/8584435" target="\_blank">8140421</a>).

## **Cellular Location**

Apical cell membrane; Multi-pass membrane protein. Basolateral cell membrane {ECO:0000250|UniProtKB:P34080}; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane; Multi-pass membrane protein. Golgi apparatus, trans-Golgi network membrane; Multi-pass membrane protein. Note=Shuttles from vesicles to the apical membrane (PubMed:15509592). Vasopressin-regulated phosphorylation is required for translocation to the apical cell membrane (PubMed:15509592). PLEKHA8/FAPP2 is required to transport AQP2 from the TGN to sites where AQP2 is phosphorylated (By similarity) {ECO:0000250|UniProtKB:P34080, ECO:0000269|PubMed:15509592}

## **Tissue Location**

Expressed in collecting tubules in kidney medulla (at protein level) (PubMed:7510718). Detected in kidney (PubMed:7510718).

# Anti-Aquaporin 2 Picoband Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Aquaporin 2 Picoband Antibody - Images

1 2 3 4 5 130KD – 70KD – 55KD – 35KD – 25KD – 15KD –

Anti- Aquaporin 2 Picoband antibody, ABO12162, Western blottingAll lanes: Anti Aquaporin 2 (ABO12162) at 0.5ug/mlLane 1: Rat Kidney Tissue Lysate at 50ugLane 2: Mouse Kidney Tissue Lysate at 50ugLane 3: HELA Whole Cell Lysate at 40ugLane 4: A549 Whole Cell Lysate at 40ugLane 5: PANC Whole Cell Lysate at 40ugPredicted bind size: 29KDObserved bind size: 50KD



Anti- Aquaporin 2 Picoband antibody, ABO12162, IHC(P)IHC(P): Mouse Kidney Tissue





Anti- Aquaporin 2 Picoband antibody, ABO12162, IHC(P)IHC(P): Rat Kidney Tissue



Anti- Aquaporin 2 Picoband antibody, ABO12162, IHC(P)IHC(P): Human Kidney Cancer Tissue Anti-Aquaporin 2 Picoband Antibody - Background

AQP2 (Aquaporin 2), also called AQUAPORIN-CD, is found in the apical cell membranes of the kidney's collecting duct principal cells and in intracellular vesicles located throughout the cell. The AQP2 gene is mapped to chromosome 12q13, very close to the site of major intrinsic protein by situ hybridization. The investigators suggested that a defect in the AOP2 gene is the basis of the autosomal form of nephrogenic diabetes insipidus. The functional expression and the limited localization suggested that AQP2 is the vasopressin-regulated water channel. Using rat kidney slices and porcine kidney cells stably expressing rat Aqp2, AQP2 trafficking can be stimulated by cAMP-independent pathways that utilize nitric oxide (NO). The NO donors sodium nitroprusside (SNP) and NONOate and the NO synthase substrate L-arginine mimicked the effect of vasopressin (VP), stimulating relocation of Aqp2 from cytoplasmic vesicles to the apical plasma membrane. SNP increased intracellular cGMP rather than cAMP, and exogenous cGMP stimulated AQP2 membrane insertion. Atrial natriuretic factor, which signals via cGMP, also stimulated AQP2 translocation. AQP2 expression in kidney connecting tubules is sufficient for survival and that AQP2 expression in collecting ducts is required to regulate body water balance. The S256L substitution in the cytoplasmic tail of the Aqp2 protein prevented phosphorylation at S256 and the subsequent accumulation of Agp2 on the apical membrane of the collecting duct principal cells.