

Anti-CCL3 Picoband Antibody
Catalog # ABO12257**Specification**

Anti-CCL3 Picoband Antibody - Product Information

Application	WB, E
Primary Accession	P10147
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for C-C motif chemokine 3(CCL3) detection. Tested with WB, ELISA in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-CCL3 Picoband Antibody - Additional Information

Gene ID 6348

Other Names

C-C motif chemokine 3, G0/G1 switch regulatory protein 19-1, Macrophage inflammatory protein 1-alpha, MIP-1-alpha, PAT 464.1, SIS-beta, Small-inducible cytokine A3, Tonsillar lymphocyte LD78 alpha protein, MIP-1-alpha(4-69), LD78-alpha(4-69), CCL3, G0S19-1, MIP1A, SCYA3

Calculated MW

10085 MW KDa

Application Details

ELISA , 0.1-0.5 µg/ml, Human, -
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Secreted.

Protein Name

C-C motif chemokine 3

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

E. coli-derived human CCL3 recombinant protein (Position: S24-A92). Human CCL3 shares 75.8% and 77.3% amino acid (aa) sequence identity with mouse and rat CCL3, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Sequence Similarities

Belongs to the intercrine beta (chemokine CC) family.

Anti-CCL3 Picoband Antibody - Protein Information

Name CCL3

Synonyms G0S19-1, MIP1A, SCYA3

Function

Monokine with inflammatory and chemokinetic properties. Binds to CCR1, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant MIP-1-alpha induces a dose- dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV).


Cellular Location


Secreted.

Anti-CCL3 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-CCL3 Picoband Antibody - Images



97KD —
58KD —
40KD —
29KD —
20KD —
14KD —

Anti- CCL3 Picoband antibody, ABO12257), Western blotting All lanes: Anti CCL3 (ABO12257) at 0.5ug/ml WB: JURKAT Whole Cell Lysate at 40ug Predicted bind size: 11KD Observed bind size: 11KD

Anti-CCL3 Picoband Antibody - Background

Macrophage inflammatory protein-1 alpha (MIP-1 alpha), also called CCL3, LD78. The cDNA for CCL3 predicts a mature peptide of 69 amino acids with a molecular mass of 7,889 daltons. CCL3 is a member of a newly identified superfamily of small inducible proteins involved in inflammatory responses, wound healing and tumorigenesis. CCL3 is a chemokine that has pro-inflammatory and stem cell inhibitory activities in vitro. It constitutes an important second signal for mast cell degranulation in the conjunctiva in vivo and consequently for acute-phase disease.