

**Anti-RANTES Picoband Antibody**  
**Catalog # ABO12259****Specification****Anti-RANTES Picoband Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">P13501</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for C-C motif chemokine 5(CCL5) detection. Tested with WB, ELISA in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-RANTES Picoband Antibody - Additional Information****Gene ID 6352****Other Names**

C-C motif chemokine 5, EoCP, Eosinophil chemotactic cytokine, SIS-delta, Small-inducible cytokine A5, T cell-specific protein P228, TCP228, T-cell-specific protein RANTES, RANTES(3-68), RANTES(4-68), CCL5, D17S136E, SCYA5

**Calculated MW**

9990 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, -<br>ELISA , 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Secreted.

**Tissue Specificity**

T-cell and macrophage specific.

**Protein Name**

C-C motif chemokine 5

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

**Immunogen**

E. coli-derived human RANTES recombinant protein (Position: Y26-S91). Human RANTES shares 83.3% amino acid (aa) sequence identity with both mouse and rat RANTES.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

**Storage**

At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Sequence Similarities**

Belongs to the intercrine beta (chemokine CC) family.

**Anti-RANTES Picoband Antibody - Protein Information****Name** CCL5**Synonyms** D17S136E, SCYA5**Function**

Chemoattractant for blood monocytes, memory T-helper cells and eosinophils. Causes the release of histamine from basophils and activates eosinophils. May activate several chemokine receptors including CCR1, CCR3, CCR4 and CCR5. One of the major HIV-suppressive factors produced by CD8+ T-cells. Recombinant RANTES protein induces a dose-dependent inhibition of different strains of HIV-1, HIV-2, and simian immunodeficiency virus (SIV). The processed form RANTES(3-68) acts as a natural chemotaxis inhibitor and is a more potent inhibitor of HIV-1-infection. The second processed form RANTES(4-68) exhibits reduced chemotactic and HIV-suppressive activity compared with RANTES(1-68) and RANTES(3-68) (PubMed:<a href="http://www.uniprot.org/citations/1380064" target="\_blank">1380064</a>, PubMed:<a href="http://www.uniprot.org/citations/15923218" target="\_blank">15923218</a>, PubMed:<a href="http://www.uniprot.org/citations/16791620" target="\_blank">16791620</a>, PubMed:<a href="http://www.uniprot.org/citations/8525373" target="\_blank">8525373</a>, PubMed:<a href="http://www.uniprot.org/citations/9516414" target="\_blank">9516414</a>). May also be an agonist of the G protein-coupled receptor GPR75, stimulating inositol trisphosphate production and calcium mobilization through its activation. Together with GPR75, may play a role in neuron survival through activation of a downstream signaling pathway involving the PI3, Akt and MAP kinases. By activating GPR75 may also play a role in insulin secretion by islet cells (PubMed:<a href="http://www.uniprot.org/citations/23979485" target="\_blank">23979485</a>).

**Cellular Location**

Secreted.

**Tissue Location**

Expressed in the follicular fluid (at protein level). T-cell and macrophage specific.

**Anti-RANTES Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-RANTES Picoband Antibody - Images**

97KD—  
58KD—  
40KD—  
29KD—  
20KD—  
14KD—

Anti-RANTES Picoband antibody, ABO12259, Western blotting All lanes: Anti RANTES (ABO12259) at 0.5ug/mlWB: HUT Whole Cell Lysate at 40ugPredicted bind size: 11KDObserved bind size: 11KD

**Anti-RANTES Picoband Antibody - Background**

Chemokine (C-C motif) ligand 5 (also CCL5) is a protein which in humans is encoded by the CCL5 gene. It is also known as RANTES. This gene is one of several chemokine genes clustered on the q-arm of chromosome 17. Chemokines form a superfamily of secreted proteins involved in immunoregulatory and inflammatory processes. The superfamily is divided into four subfamilies based on the arrangement of the N-terminal cysteine residues of the mature peptide. This chemokine, a member of the CC subfamily, functions as a chemoattractant for blood monocytes, memory T helper cells and eosinophils. It causes the release of histamine from basophils and activates eosinophils. This cytokine is one of the major HIV-suppressive factors produced by CD8+ cells. It functions as one of the natural ligands for the chemokine receptor chemokine (C-C motif) receptor 5 (CCR5), and it suppresses in vitro replication of the R5 strains of HIV-1, which use CCR5 as a coreceptor. Alternative splicing results in multiple transcript variants that encode different isoforms.