

Anti-HDAC6 Picoband Antibody
Catalog # ABO12314**Specification**

Anti-HDAC6 Picoband Antibody - Product Information

Application	WB
Primary Accession	Q9UBN7
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Histone deacetylase 6(HDAC6) detection. Tested with WB in Human;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-HDAC6 Picoband Antibody - Additional Information

Gene ID 10013

Other Names

Histone deacetylase 6, HD6, 3.5.1.98, HDAC6, KIAA0901

Calculated MW

131419 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Rat

Subcellular Localization

Nucleus. Cytoplasm. Perikaryon . Cell projection, dendrite . Cell projection, axon . It is mainly cytoplasmic, where it is associated with microtubules.

Protein Name

Histone deacetylase 6

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human HDAC6 (137-169aa EKEELMLVHSLEYIDLMETTQYMNEGELRVLAD), different from the related mouse sequence by one amino acid.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-HDAC6 Picoband Antibody - Protein Information

Name HDAC6 {ECO:0000303|PubMed:10220385, ECO:0000312|HGNC:HGNC:14064}

Function

Deacetylates a wide range of non-histone substrates (PubMed:12024216, PubMed:18606987, PubMed:20308065, PubMed:24882211, PubMed:26246421, PubMed:30538141, PubMed:31857589, PubMed:30770470, PubMed:38534334, PubMed:39567688). Plays a central role in microtubule- dependent cell motility by mediating deacetylation of tubulin (PubMed:12024216, PubMed:20308065, PubMed:26246421). Required for cilia disassembly via deacetylation of alpha-tubulin (PubMed:17604723, PubMed:26246421). Alpha-tubulin deacetylation results in destabilization of dynamic microtubules (By similarity). Promotes deacetylation of CTTN, leading to actin polymerization, promotion of autophagosome-lysosome fusion and completion of autophagy (PubMed:30538141). Deacetylates SQSTM1 (PubMed:31857589). Deacetylates peroxiredoxins PRDX1 and PRDX2, decreasing their reducing activity (PubMed:18606987). Deacetylates antiviral protein RIGI in the presence of viral mRNAs which is required for viral RNA detection by RIGI (By similarity). Sequentially deacetylates and polyubiquitinates DNA mismatch repair protein MSH2 which leads to MSH2 degradation, reducing cellular sensitivity to DNA-damaging agents and decreasing cellular DNA mismatch repair activities (PubMed:24882211). Deacetylates DNA mismatch repair protein MLH1 which prevents recruitment of the MutL alpha complex (formed by the MLH1-PMS2 heterodimer) to the MutS alpha complex (formed by the MSH2-MSH6 heterodimer), leading to tolerance of DNA damage (PubMed:30770470). Deacetylates RHOT1/MIRO1 which blocks mitochondrial transport and mediates axon growth inhibition (By similarity). Deacetylates transcription factor SP1 which leads to increased expression of ENG, positively regulating angiogenesis (PubMed:38534334). Deacetylates KHDRBS1/SAM68 which regulates alternative splicing by inhibiting the inclusion of CD44 alternate exons (PubMed:26080397). Acts as a valine sensor by binding to valine through the primate-specific SE14 repeat region (PubMed:<a

[39567688](http://www.uniprot.org/citations/39567688)). In valine deprivation conditions, translocates from the cytoplasm to the nucleus where it deacetylates TET2 which promotes TET2-dependent DNA demethylation, leading to DNA damage (PubMed:[39567688](http://www.uniprot.org/citations/39567688)). Promotes odontoblast differentiation following IPO7-mediated nuclear import and subsequent repression of RUNX2 expression (By similarity). In addition to its protein deacetylase activity, plays a key role in the degradation of misfolded proteins: when misfolded proteins are too abundant to be degraded by the chaperone refolding system and the ubiquitin-proteasome, mediates the transport of misfolded proteins to a cytoplasmic juxtanuclear structure called aggresome (PubMed:[17846173](http://www.uniprot.org/citations/17846173)). Probably acts as an adapter that recognizes polyubiquitinated misfolded proteins and targets them to the aggresome, facilitating their clearance by autophagy (PubMed:[17846173](http://www.uniprot.org/citations/17846173)). Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer (PubMed:[24413532](http://www.uniprot.org/citations/24413532)).

Cellular Location


Cytoplasm. Cytoplasm, cytoskeleton. Nucleus. Perikaryon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, axon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, cilium. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body Note=Mainly cytoplasmic where it is associated with microtubules (PubMed:12024216). Can shuttle between the cytoplasm and the nucleus (PubMed:39567688). Retained in the cytoplasm by binding to valine via the primate-specific SE14 repeat region while valine deprivation induces nuclear localization (PubMed:39567688). Found exclusively in the cytoplasm in proliferative cells with a fraction found in the nucleus during differentiation (By similarity). May translocate to the nucleus following DNA damage (PubMed:30770470) {ECO:0000250|UniProtKB:Q9Z2V5, ECO:0000269|PubMed:12024216, ECO:0000269|PubMed:30770470, ECO:0000269|PubMed:39567688}


Anti-HDAC6 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-HDAC6 Picoband Antibody - Images





Anti- HDAC6 Picoband antibody, ABO12314, Western blotting All lanes: Anti HDAC6 (ABO12314) at 0.5ug/ml Lane 1: Rat Skeletal Muscle Tissue Lysate at 50ug Lane 2: COLO320 Whole Cell Lysate at 40ug Predicted bind size: 160KD Observed bind size: 160KD

Anti-HDAC6 Picoband Antibody - Background

HDAC6, also called KIAA0901, is a member belongs to class II of the histone deacetylase/acuc/apha family of proteins that is an enzyme that in humans is encoded by the HDAC6 gene. The HDAC6 gene is mapped to chromosome Xp11.23. HDAC6 contains an internal duplication of two catalytic domains which appear to function independently of each other. The protein possesses histone deacetylase activity and represses transcription. HDAC6 functions as a tubulin deacetylase. And it is localized exclusively in the cytoplasm, where it associates with microtubules and localizes with the microtubule motor complex. HDAC6 could bind both polyubiquitinated misfolded proteins and dynein motors, thereby recruiting misfolded protein cargo to dynein motors for transport to aggresomes. Furthermore, expression of HDAC6 was sufficient to rescue degeneration associated with UPS dysfunction in vivo in an autophagy-dependent manner. HDAC6 is a central component of the stress response that regulates SG formation and potentially contributes to control of RNA metabolism and translation.