

Anti-Hex Picoband Antibody

Catalog # ABO12317

Specification

Anti-Hex Picoband Antibody - Product Information

ApplicationWBPrimary Accession003014HostRabbitReactivityHuman, Mouse, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Hematopoietically-expressed homeobox protein Hhex(HHEX)detection. Tested with WB in Human;Mouse; Rat.

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Hex Picoband Antibody - Additional Information

Gene ID 3087

Other Names Hematopoietically-expressed homeobox protein HHEX, Homeobox protein HEX, Homeobox protein PRH, HHEX, HEX, PRH, PRHX

Calculated MW 30022 MW KDa

Application Details Western blot, 0.1-0.5 μg/ml, Human, Mouse, Rat

Subcellular Localization Nucleus .

Tissue Specificity Liver and promyelocytic leukemia cell line HL- 60. .

Protein Name Hematopoietically-expressed homeobox protein Hhex

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence in the middle region of human Hex(146-180aa NDQTIELEKKFETQKYLSPPERKRLAKMLQLSERQ), different from the related mouse sequence by one amino acid.



Purification Immunogen affinity purified.

Cross Reactivity No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-Hex Picoband Antibody - Protein Information

Name HHEX

Synonyms HEX, PRH, PRHX

Function

Recognizes the DNA sequence 5'-ATTAA-3' (By similarity). Transcriptional repressor (By similarity). Activator of WNT-mediated transcription in conjunction with CTNNB1 (PubMed:20028982). Establishes anterior identity at two levels; acts early to enhance canonical WNT- signaling by repressing expression of TLE4, and acts later to inhibit NODAL-signaling by directly targeting NODAL (By similarity). Inhibits EIF4E-mediated mRNA nuclear export (PubMed:12554669). May play a role in hematopoietic differentiation (PubMed:12554669). May play a role in hematopoietic differentiation (PubMed:8096636).

Cellular Location Nucleus {ECO:0000250|UniProtKB:P43120}. Nucleus, nuclear body. Cytoplasm

Tissue Location Liver and promyelocytic leukemia cell line HL-60.

Anti-Hex Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-Hex Picoband Antibody - Images





Anti- Hex Picoband antibody, ABO12317, Western blottingAll lanes: Anti Hex (ABO12317) at 0.5ug/mlLane 1: Rat Liver Tissue Lysate at 50ugLane 2: Mouse Liver Tissue Lysate at 50ugLane 3: HEPG Whole Cell Lysate at 40ugPredicted bind size: 47KDObserved bind size: 47KD

Anti-Hex Picoband Antibody - Background

Hematopoietically-expressed homeobox protein HHEX is a protein that in humans is encoded by the HHEX gene. Homeobox genes are members of a family of transcription factors that regulate tissue development in many different organisms. Hromas et al. (1993) set out to identify homeobox genes that might play a role in hematopoiesis. And using somatic cell hybrid analysis, they mapped the HHEX gene to chromosome 10, where the HOX11 gene is located. Homeobox genes are involved in neoplastic transformation of both epithelial and hemopoietic tissues. The divergent homeobox gene HEX is expressed in the anterior visceral endoderm during early mouse development and in some adult tissues of endodermal origin, including liver and thyroid. D'Elia et al.'s findings suggested that regulation of HEX entry in the nucleus of thyrocytes may represent a critical step during human thyroid tumorigenesis.