

**Anti-KChIP2 Picoband Antibody**  
**Catalog # ABO12338****Specification**

---

**Anti-KChIP2 Picoband Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">Q9NS61</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Kv channel-interacting protein 2(KCNIP2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-KChIP2 Picoband Antibody - Additional Information**

**Gene ID** 30819

**Other Names**

Kv channel-interacting protein 2, KChIP2, A-type potassium channel modulatory protein 2, Cardiac voltage-gated potassium channel modulatory subunit, Potassium channel-interacting protein 2, KCNIP2, KCHIP2

**Calculated MW**

30907 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat  
Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

**Subcellular Localization**

Isoform 1: Cell membrane ; Lipid-anchor . Detected on lipid rafts (By similarity). .

**Tissue Specificity**

Expressed in brain. Colocalizes with KCND2 in excitatory neurons including cortical and hippocampal CA1 pyramidal cells. Isoform 3 is expressed in heart and in umbilical vein endothelial cells. Not expressed in fetal heart. .

**Protein Name**

Kv channel-interacting protein 2

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

**Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human KChIP2 (78-112aa DEFELSTVCHRPEGLEQLQEQTkFTRKELQVLYR), different from the related mouse and rat sequences by one amino acid.

#### **Purification**

Immunogen affinity purified.

#### **Cross Reactivity**

No cross reactivity with other proteins

#### **Storage**

**At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.**

### **Anti-KChIP2 Picoband Antibody - Protein Information**

**Name** KCNIP2 ([HGNC:15522](#))

#### **Function**

Regulatory subunit of Kv4/D (Shal)-type voltage-gated rapidly inactivating A-type potassium channels (PubMed: [10676964](http://www.uniprot.org/citations/10676964), PubMed: [11287421](http://www.uniprot.org/citations/11287421), PubMed: [11684073](http://www.uniprot.org/citations/11684073), PubMed: [12297301](http://www.uniprot.org/citations/12297301), PubMed: [14623880](http://www.uniprot.org/citations/14623880), PubMed: [34997220](http://www.uniprot.org/citations/34997220)). Modulates channel density, inactivation kinetics and rate of recovery from inactivation in a calcium-dependent and isoform-specific manner (PubMed: [10676964](http://www.uniprot.org/citations/10676964), PubMed: [11287421](http://www.uniprot.org/citations/11287421), PubMed: [11684073](http://www.uniprot.org/citations/11684073), PubMed: [12297301](http://www.uniprot.org/citations/12297301), PubMed: [14623880](http://www.uniprot.org/citations/14623880), PubMed: [34997220](http://www.uniprot.org/citations/34997220)). Involved in KCND2 and KCND3 trafficking to the cell surface (PubMed: [12829703](http://www.uniprot.org/citations/12829703)). May be required for the expression of I(To) currents in the heart (By similarity).

#### **Cellular Location**

[Isoform 1]: Cell membrane {ECO:0000250|UniProtKB:Q9JM59}; Lipid-anchor {ECO:0000250|UniProtKB:Q9JM59}. Note=Detected on lipid rafts (By similarity). {ECO:0000250|UniProtKB:Q9JM59} [Isoform 6]: Cell membrane {ECO:0000250|UniProtKB:Q9JM59}; Lipid-anchor {ECO:0000250|UniProtKB:Q9JM59}

#### **Tissue Location**


Expressed in brain. Colocalizes with KCND2 in excitatory neurons including cortical and hippocampal CA1 pyramidal cells. Isoform 3 is expressed in heart and in umbilical vein endothelial cells. Not expressed in fetal heart

### **Anti-KChIP2 Picoband Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


#### Anti-KCHIP2 Picoband Antibody - Images




Anti- KCNIP2 Picoband antibody, ABO12338, Western blotting All lanes: Anti KCNIP2 (ABO12338) at 0.5ug/ml  
Lane 1: Rat Brain Tissue Lysate at 50ug  
Lane 2: Rat Cardiac Muscle Tissue Lysate at 50ug  
Lane 3: Mouse Cardiac Muscle Tissue Lysate at 50ug  
Lane 4: 22RV1 Whole Cell Lysate at 40ug  
Predicted bind size: 31KD  
Observed bind size: 31KD



Anti- KCNIP2 Picoband antibody, ABO12338, IHC(P) IHC(P): Mouse Brain Tissue



Anti- KCNIP2 Picoband antibody, ABO12338,IHC(P)IHC(P): Human Glioma Tissue



Anti- KCNIP2 Picoband antibody, ABO12338,IHC(P)IHC(P): Rat Brain Tissue

#### **Anti-KChIP2 Picoband Antibody - Background**

Kv channel-interacting protein 2 also known as KChIP2 is a protein that in humans is encoded by the KCNIP2 gene. This gene encodes a member of the family of voltage-gated potassium (Kv) channel-interacting proteins, which belong to the recoverin branch of the EF-hand superfamily. Members of the KCNIP family are small calcium binding proteins. They all have EF-hand-like domains, and differ from each other in the N-terminus. And they are integral subunit components of native Kv4 channel complexes. They may regulate A-type currents, and hence neuronal excitability, in response to changes in intracellular calcium. Alternative splicing results in multiple transcript variant encoding different isoforms.