

## **Anti-MAPK6 Picoband Antibody**

**Catalog # ABO12352** 

## **Specification**

# **Anti-MAPK6 Picoband Antibody - Product Information**

Application WB, IHC-P
Primary Accession Q16659
Host Rabbit

Reactivity Human, Mouse, Rat

Clonality Polyclonal Lyophilized

**Description** 

Rabbit IgG polyclonal antibody for Mitogen-activated protein kinase 6(MAPK6) detection. Tested with WB, IHC-P in Human; Mouse; Rat.

### Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

## **Anti-MAPK6 Picoband Antibody - Additional Information**

**Gene ID 5597** 

#### **Other Names**

Mitogen-activated protein kinase 6, MAP kinase 6, MAPK 6, 2.7.11.24, Extracellular signal-regulated kinase 3, ERK-3, MAP kinase isoform p97, p97-MAPK, MAPK6, ERK3, PRKM6

## Calculated MW 82681 MW KDa

### **Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μg/ml, Human, By Heat<br/>br>Western blot, 0.1-0.5 μg/ml, Human, Mouse, Rat<br/>br>

### **Subcellular Localization**

Cytoplasm . Nucleus . Translocates to the cytoplasm following interaction with MAPKAPK5. .

### **Tissue Specificity**

Highest expression in the skeletal muscle, followed by the brain. Also found in heart, placenta, lung, liver, pancreas, kidney and skin fibroblasts.

### **Protein Name**

Mitogen-activated protein kinase 6

### Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

### **Immunogen**

E.coli-derived human MAPK6 recombinant protein (Position: E520-N721). Human MAPK6 shares 86.2% and 82.8% amino acid (aa) sequence identity with mouse and rat MAPK6, respectively.



Purification Immunogen affinity purified.

**Cross Reactivity**No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

## **Anti-MAPK6 Picoband Antibody - Protein Information**

Name MAPK6

Synonyms ERK3, PRKM6

### **Function**

Atypical MAPK protein. Phosphorylates microtubule-associated protein 2 (MAP2) and MAPKAPK5. The precise role of the complex formed with MAPKAPK5 is still unclear, but the complex follows a complex set of phosphorylation events: upon interaction with atypical MAPKAPK5, ERK3/MAPK6 is phosphorylated at Ser-189 and then mediates phosphorylation and activation of MAPKAPK5, which in turn phosphorylates ERK3/MAPK6. May promote entry in the cell cycle (By similarity).

### **Cellular Location**

Cytoplasm. Nucleus. Note=Translocates to the cytoplasm following interaction with MAPKAPK5

## **Tissue Location**

Highest expression in the skeletal muscle, followed by the brain. Also found in heart, placenta, lung, liver, pancreas, kidney and skin fibroblasts


## **Anti-MAPK6 Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## **Anti-MAPK6 Picoband Antibody - Images**





Anti- MAPK6 Picoband antibody, ABO12352, Western blottingAll lanes: Anti MAPK6 (ABO12352) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Rat Skeletal Muscle Tissue Lysate at 50ugLane 3: PANC Whole Cell Lysate at 40ugLane 4: NIH3T3 Whole Cell Lysate at 40ugPredicted bind size: 68KDObserved bind size: 100KD



Anti- MAPK6 Picoband antibody, ABO12352, IHC(P)IHC(P): Human Lung Cancer Tissue

## Anti-MAPK6 Picoband Antibody - Background

Mitogen-activated protein kinase 6 is an enzyme that in humans is encoded by the MAPK6 gene. It is mapped to 15q21. The protein encoded by this gene is a member of the Ser/Thr protein kinase family, and is most closely related to mitogen-activated protein kinases (MAP kinases). MAP kinases also known as extracellular signal-regulated kinases (ERKs), are activated through protein phosphorylation cascades and act as integration points for multiple biochemical signals. This kinase is localized in the nucleus, and has been reported to be activated in fibroblastsupon treatment with serum or phorbol esters.