

Anti-IGFBP5 Picoband Antibody
Catalog # ABO12397**Specification**

Anti-IGFBP5 Picoband Antibody - Product Information

Application	WB, E
Primary Accession	P24593
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Insulin-like growth factor-binding protein 5(IGFBP5) detection. Tested with WB, ELISA in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-IGFBP5 Picoband Antibody - Additional Information

Gene ID 3488

Other Names

Insulin-like growth factor-binding protein 5, IBP-5, IGF-binding protein 5, IGFBP-5, IGFBP5, IBP5

Calculated MW

30570 MW KDa

Application Details

ELISA , 0.1-0.5 µg/ml, Human, -
Western blot, 0.1-0.5 µg/ml, Human

Subcellular Localization

Secreted.

Tissue Specificity

Osteosarcoma, and at lower levels in liver, kidney and brain.

Protein Name

Insulin-like growth factor-binding protein 5

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the N-terminus of human IGFBP5 (76-114aa QGLRCLPRQDEEKPLHALLHGRGVCLNEKSYREQVKIER), different from the related mouse and rat sequences by two amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-IGFBP5 Picoband Antibody - Protein Information

Name IGFBP5

Synonyms IBP5

Function

Multifunctional protein that plays a critical role in regulating the availability of IGFs to their receptors and thereby regulates IGF-mediated cellular processes including proliferation, differentiation, and apoptosis in a cell-type specific manner (PubMed:18930415, PubMed:7683690). Increases the cell proliferation of osteoblasts, intestinal smooth muscle cells and neuroblastoma cells. Enhances adhesion and survival of epithelial cells but decreases adhesion of mesenchymal cells (By similarity). Once secreted, acts as a major mediator of mTORC1-dependent feedback inhibition of IGF1 signaling (By similarity). Also plays a role in the induction of extracellular matrix (ECM) production and deposition independently of its nuclear translocation and binding to IGFs (PubMed:20345844, PubMed:26103640). Acts itself as a growth factor that can act independently of IGFs to regulate bone formation. Acts as a ligand for the ROR1 receptor which triggers formation of ROR1/HER2 heterodimer to enhance CREB oncogenic signaling (PubMed:36949068).

Cellular Location

Secreted. Cytoplasm. Nucleus

Tissue Location


Osteosarcoma, and at lower levels in liver, kidney and brain

Anti-IGFBP5 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-IGFBP5 Picoband Antibody - Images



Anti- IGFBP5 Picoband antibody, ABO12397, Western blotting All lanes: Anti IGFBP5 (ABO12397) at 0.5ug/ml
Lane 1: U20S Whole Cell Lysate at 40ug
Lane 2: HELA Whole Cell Lysate at 40ug
Predicted bind size: 31KD
Observed bind size: 23KD

Anti-IGFBP5 Picoband Antibody - Background

Insulin-like growth factor-binding protein 5 is a protein that in humans is encoded by the IGFBP5 gene. The expression of IGFBP5 by stable transfection and adenovirus-mediated infection is inhibitory to growth in 2 human breast cancer cell lines. IGFBP5 expression leads to G2/M cell cycle arrest and apoptosis. Stable expression of IGFBP5 in the breast cancer cell lines also inhibits the formation and growth of tumors following injection in athymic mice. It is concluded that IGFBP5 is a growth inhibitor and proapoptotic agent in breast cancer cells. Additionally, IGFBP-5 is expressed by fibroblasts, myoblasts and osteoblasts, making it the predominant IGFBP found in bone extracts. It has a strong affinity for hydroxyapatite, allowing it to bind to bone cells. When bound to extracellular matrix, IGFBP-5 is protected from proteolysis and potentiates IGF activity, but when it is soluble, IGFBP-5 is cleaved to a biologically inactive 21 kDa fragment (1, 2).