

## **Anti-OMG Picoband Antibody**

Catalog # ABO12416

## **Specification**

# **Anti-OMG Picoband Antibody - Product Information**

Application WB
Primary Accession P23515
Host Rabbit

Reactivity Human, Mouse, Rat

Clonality Polyclonal Lyophilized

**Description** 

Rabbit IgG polyclonal antibody for Oligodendrocyte-myelin glycoprotein(OMG) detection. Tested with WB in Human; Mouse; Rat.

## Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

## **Anti-OMG Picoband Antibody - Additional Information**

**Gene ID 4974** 

**Other Names** 

Oligodendrocyte-myelin glycoprotein, OMG, OMGP

Calculated MW 49608 MW KDa

**Application Details** 

Western blot, 0.1-0.5 μg/ml, Human, Mouse, Rat<br/>
br>

**Subcellular Localization** 

Cell membrane; Lipid-anchor, GPI-anchor.

**Tissue Specificity** 

Oligodendrocytes and myelin of the central nervous system.

**Protein Name** 

Oligodendrocyte-myelin glycoprotein

**Contents** 

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

#### **Immunogen**

A synthetic peptide corresponding to a sequence at the N-terminus of human OMG (64-99aa YNHFTDLHNQLTQYTNLRTLDISNNRLESLPAHLPR), different from the related mouse sequence by two amino acids.

#### **Purification**



Immunogen affinity purified.

**Cross Reactivity** 

No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

## **Anti-OMG Picoband Antibody - Protein Information**

Name OMG

Synonyms OMGP

#### **Function**

Cell adhesion molecule contributing to the interactive process required for myelination in the central nervous system.

#### **Cellular Location**

Cell membrane; Lipid-anchor, GPI-anchor.

#### **Tissue Location**

Oligodendrocytes and myelin of the central nervous system


#### **Anti-OMG Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **Anti-OMG Picoband Antibody - Images**





Anti- OMG Picoband antibody, ABO12416, Western blottingAll lanes: Anti OMG (ABO12416) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Mouse Brain Tissue Lysate at 50ugLane 3: U87 Whole Cell Lysate at 40ugPredicted bind size: 50KDObserved bind size: 50KD

## **Anti-OMG Picoband Antibody - Background**

Oligodendrocyte-myelin glycoprotein is a protein that in humans is encoded by the OMG gene. This gene is mapped to 17q11-q12. OMG, like neurofibromin (NF1), has growth suppressive effects and downregulates mitogenic signaling pathways closely related to those influenced by neurofibromin. Overexpression of OMG alters mitogenic signaling in fibroblasts. Cells overexpressing OMG grow more slowly in serum compared to controls and show a partial G1 block upon cell cycle analysis.