

**Anti-SDC1/Syndecan-1 Antibody**  
**Catalog # ABO12429****Specification****Anti-SDC1/Syndecan-1 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P18827</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Syndecan-1( SDC1) detection. Tested with WB in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-SDC1/Syndecan-1 Antibody - Additional Information****Gene ID** 6382**Other Names**

Syndecan-1, SYND1, CD138, SDC1, SDC

**Calculated MW**

32462 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human<br>

**Subcellular Localization**

Membrane; Single-pass type I membrane protein. Secreted. Shedding of the ectodomain produces a soluble form.

**Protein Name**

Syndecan-1

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

**Immunogen**

A synthetic peptide corresponding to a sequence at the C-terminus of human Syndecan-1 (218-251aa ENTAVVAVEPDRRNQSPV рDQGATGASQGLLDRKE), different from the related mouse sequence by six amino acids, and from the related rat sequence by four amino acids.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins.

Storage

At -20°C for one year. After r° Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

### Anti-SDC1/Syndecan-1 Antibody - Protein Information

Name SDC1 ([HGNC:10658](#))

Synonyms SDC

#### Function

Cell surface proteoglycan that contains both heparan sulfate and chondroitin sulfate and that links the cytoskeleton to the interstitial matrix (By similarity). Regulates exosome biogenesis in concert with SDCBP and PDCD6IP (PubMed:<a href="http://www.uniprot.org/citations/22660413" target="\_blank">22660413</a>). Able to induce its own expression in dental mesenchymal cells and also in the neighboring dental epithelial cells via an MSX1-mediated pathway (By similarity).

#### Cellular Location

Membrane; Single-pass type I membrane protein. Secreted Secreted, extracellular exosome  
Note=Shedding of the ectodomain produces a soluble form

#### Tissue Location


Detected in placenta (at protein level) (PubMed:32337544). Detected in fibroblasts (at protein level) (PubMed:36213313).

### Anti-SDC1/Syndecan-1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-SDC1/Syndecan-1 Antibody - Images



Anti- Syndecan-1 Picoband antibody, ABO12429, Western blottingAll lanes: Anti Syndecan-1 (ABO12429) at 0.5ug/mlLane 1: HT1080 Whole Cell Lysate at 40ugLane 2: MCF-7 Whole Cell Lysate at 40ugLane 3: 293T Whole Cell Lysate at 40ugLane 4: HEPG2 Whole Cell Lysate at 40ugLane 5: HEGLA Whole Cell Lysate at 40ugPredicted bind size: 32KDObserved bind size: 50KD

#### Anti-SDC1/Syndecan-1 Antibody - Background

Syndecan 1 is a protein which in humans is encoded by the SDC1 gene. The protein encoded by this gene is a transmembrane (type I) heparan sulfate proteoglycan and is a member of the syndecan proteoglycan family. The syndecans mediate cell binding, cell signaling, and cytoskeletal organization and syndecan receptors are required for internalization of the HIV-1 tat protein. The syndecan-1 protein functions as an integral membrane protein and participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. Altered syndecan-1 expression has been detected in several different tumor types. While several transcript variants may exist for this gene, the full-length natures of only two have been described to date. These two represent the major variants of this gene and encode the same protein.