

Anti-PPT1 Picoband Antibody

Catalog # ABO12467

Specification

Anti-PPT1 Picoband Antibody - Product Information

ApplicationWB, IHC-PPrimary AccessionP50897HostRabbitReactivityHuman, RatClonalityPolyclonalFormatLyophilizedDescriptionRabbit IgG polyclonal antibody for Palmitoyl-protein thioesterase 1(PPT1) detection. Tested withWB, IHC-P in Human;Rat.WB, IHC-P

Reconstitution Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-PPT1 Picoband Antibody - Additional Information

Gene ID 5538

Other Names Palmitoyl-protein thioesterase 1, PPT-1, 3.1.2.22, Palmitoyl-protein hydrolase 1, PPT1, CLN1 {ECO:0000303|PubMed:19941651}, PPT

Calculated MW 34193 MW KDa

Application Details Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μg/ml, Human, By Heat

Western blot, 0.1-0.5 μg/ml, Human, Rat

Subcellular Localization Lysosome.

Protein Name Palmitoyl-protein thioesterase 1

Contents Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human PPT1 (191-224aa KEDVYRNHSIFLADINQERGINESYKKNLMALKK), different from the related mouse and rat sequences by four amino acids.

Purification Immunogen affinity purified.



Cross Reactivity No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.

Anti-PPT1 Picoband Antibody - Protein Information

Name PPT1

Synonyms CLN1 {ECO:0000303|PubMed:19941651}, PPT

Function

Has thioesterase activity against fatty acid thioesters with 14 -18 carbons, including palmitoyl-CoA, S-palmitoyl-N- acetylcysteamine, and palmitoylated proteins (PubMed:12855696, PubMed:26731412, PubMed:8816748). In contrast to PPT2, PPT1 can hydrolyze palmitoylated proteins and palmitoylcysteine (PubMed:12855696). In contrast to PPT2, PPT1 can hydrolyze palmitoylated proteins and palmitoylcysteine (PubMed:12855696).

Cellular Location Lysosome. Secreted Golgi apparatus. Endoplasmic reticulum

Anti-PPT1 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

Anti-PPT1 Picoband Antibody - Images



Figure 1. Western blot analysis of PPT1 using anti- PPT1 antibody (ABO12467). Electrophoresis



was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates, Lane 3: rat liver tissue lysates, Lane 4: mouse liver tissue lysates, Lane 5: HEPG2 whole Cell lysates, Lane 6: 293T whole cell lysates, Lane 7: MCF-7 whole cell lysates. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti- PPT1 antigen affinity purified polyclonal antibody (Catalog # ABO12467) at 0.5 $\hat{1}_{4}$ g/mL overnight at 4 \hat{A} °C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for PPT1.



Figure 2. IHC analysis of PPT1 using anti- PPT1 antibody (ABO12467).PPT1 was detected in paraffin-embedded section of human intestinal cancer tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with $11\frac{1}{4}$ g/ml rabbit anti- PPT1 Antibody (ABO12467) overnight at $4\hat{A}^\circ$ C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at $37\hat{A}^\circ$ C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC) with DAB as the chromogen.

Anti-PPT1 Picoband Antibody - Background

Palmitoyl-protein thioesterase 1 (PPT-1), also known as palmitoyl-protein hydrolase 1, is an enzyme that in humans is encoded by the PPT1 gene. PPT-1 is a member of the palmitoyl protein thioesterase family. The protein encoded by this gene is a small glycoprotein involved in the catabolism of lipid-modified proteins during lysosomal degradation. The encoded enzyme removes thioester-linked fatty acyl groups such as palmitate from cysteine residues. Defects in this gene are a cause of infantile neuronal ceroid lipofuscinosis 1 (CLN1, or INCL) and neuronal ceroid lipofuscinosis 4 (CLN4). Two transcript variants encoding different isoforms have been found for this gene.