

**Anti-POLH Picoband Antibody**  
**Catalog # ABO12499****Specification**

---

**Anti-POLH Picoband Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q9Y253</a>
Host	Rabbit
Reactivity	Human, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for DNA polymerase eta(POLH) detection. Tested with WB in Human;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-POLH Picoband Antibody - Additional Information**

**Gene ID** 5429

**Other Names**

DNA polymerase eta, 2.7.7.7, RAD30 homolog A, Xeroderma pigmentosum variant type protein, POLH, RAD30, RAD30A, XPV

**Calculated MW**

78413 MW KDa

**Application Details**

Western blot, 0.1-0.5 µg/ml, Human, Rat<br>

**Subcellular Localization**

Nucleus . Accumulates at replication forks after DNA damage.

**Protein Name**

DNA polymerase eta

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immunogen**

E.coli-derived human POLH recombinant protein (Position: A157-R361). Human POLH shares 93.6% amino acid (aa) sequence identity with mouse POLH.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins.

Storage

**At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time.Avoid repeated freezing and thawing.**

## Anti-POLH Picoband Antibody - Protein Information

**Name** POLH

**Synonyms** RAD30, RAD30A, XPV

### Function

DNA polymerase specifically involved in the DNA repair by translesion synthesis (TLS) (PubMed:<a href="http://www.uniprot.org/citations/10385124" target="\_blank">10385124</a>, PubMed:<a href="http://www.uniprot.org/citations/11743006" target="\_blank">11743006</a>, PubMed:<a href="http://www.uniprot.org/citations/24449906" target="\_blank">24449906</a>, PubMed:<a href="http://www.uniprot.org/citations/24553286" target="\_blank">24553286</a>, PubMed:<a href="http://www.uniprot.org/citations/16357261" target="\_blank">16357261</a>). Due to low processivity on both damaged and normal DNA, cooperates with the heterotetrameric (REV3L, REV7, POLD2 and POLD3) POLZ complex for complete bypass of DNA lesions. Inserts one or 2 nucleotide(s) opposite the lesion, the primer is further extended by the tetrameric POLZ complex. In the case of 1,2-intrastrand d(GpG)-cisplatin cross-link, inserts dCTP opposite the 3' guanine (PubMed:<a href="http://www.uniprot.org/citations/24449906" target="\_blank">24449906</a>). Particularly important for the repair of UV-induced pyrimidine dimers (PubMed:<a href="http://www.uniprot.org/citations/10385124" target="\_blank">10385124</a>, PubMed:<a href="http://www.uniprot.org/citations/11743006" target="\_blank">11743006</a>). Although inserts the correct base, may cause base transitions and transversions depending upon the context. May play a role in hypermutation at immunoglobulin genes (PubMed:<a href="http://www.uniprot.org/citations/11376341" target="\_blank">11376341</a>, PubMed:<a href="http://www.uniprot.org/citations/14734526" target="\_blank">14734526</a>). Forms a Schiff base with 5'- deoxyribose phosphate at abasic sites, but does not have any lyase activity, preventing the release of the 5'-deoxyribose phosphate (5'- dRP) residue. This covalent trapping of the enzyme by the 5'-dRP residue inhibits its DNA synthetic activity during base excision repair, thereby avoiding high incidence of mutagenesis (PubMed:<a href="http://www.uniprot.org/citations/14630940" target="\_blank">14630940</a>). Targets POLI to replication foci (PubMed:<a href="http://www.uniprot.org/citations/12606586" target="\_blank">12606586</a>).

### Cellular Location

Nucleus. Note=Binding to ubiquitinated PCNA mediates colocalization to replication foci during DNA replication and persists at sites of stalled replication forks following UV irradiation (PubMed:12606586, PubMed:16357261, PubMed:24553286). After UV irradiation, recruited to DNA damage sites within 1 hour, to a maximum of about 80%; this recruitment may not be not restricted to cells active in DNA replication (PubMed:22801543). Colocalizes with TRAP1 to nuclear foci (PubMed:24553286).


## Anti-POLH Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Anti-POLH Picoband Antibody - Images



Anti- POLH Picoband antibody, ABO12499, Western blotting All lanes: Anti POLH (ABO12499) at 0.5ug/ml  
Lane 1: Rat Liver Tissue Lysate at 50ug  
Lane 2: HELA Whole Cell Lysate at 40ug  
Lane 3: SW620 Whole Cell Lysate at 40ug  
Predicted bind size: 78KD  
Observed bind size: 78KD

### Anti-POLH Picoband Antibody - Background

DNA polymerase eta (Pol  $\eta$ ), is a protein that in humans is encoded by the POLH gene. This gene encodes a member of the Y family of specialized DNA polymerases. It copies undamaged DNA with a lower fidelity than other DNA-directed polymerases. However, it accurately replicates UV-damaged DNA; when thymine dimers are present, this polymerase inserts the complementary nucleotides in the newly synthesized DNA, thereby bypassing the lesion and suppressing the mutagenic effect of UV-induced DNA damage. This polymerase is thought to be involved in hypermutation during immunoglobulin class switch recombination. Mutations in this gene result in XPV, a variant type of xeroderma pigmentosum. Several transcript variants encoding different isoforms have been found for this gene.