

Anti-UNC5C Picoband Antibody
Catalog # ABO12527**Specification**

Anti-UNC5C Picoband Antibody - Product Information

Application	WB
Primary Accession	O95185
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Netrin receptor UNC5C(UNC5C) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-UNC5C Picoband Antibody - Additional Information

Gene ID 8633

Other Names

Netrin receptor UNC5C, Protein unc-5 homolog 3, Protein unc-5 homolog C, UNC5C, UNC5H3

Calculated MW

103146 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cell membrane ; Single-pass type I membrane protein . Cell junction, synapse, synaptosome .

Tissue Specificity

Mainly expressed in brain. Also expressed in kidney. Not expressed in developing or adult lung. .

Protein Name

Netrin receptor UNC5C

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human UNC5C (894-930aa DLWEAQNFDPGNLSMLAAVLEEMGRHETTVVSLAAEGQ), identical to the related mouse and rat sequences.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-UNC5C Picoband Antibody - Protein Information

Name UNC5C

Synonyms UNC5H3

Function

Receptor for netrin required for axon guidance (By similarity). Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding (By similarity). NTN1/Netrin-1 binding might cause dissociation of UNC5C from polymerized TUBB3 in microtubules and thereby lead to increased microtubule dynamics and axon repulsion (PubMed:28483977). Axon repulsion in growth cones may also be caused by its association with DCC that may trigger signaling for repulsion (By similarity). Might also collaborate with DSCAM in NTN1-mediated axon repulsion independently of DCC (By similarity). Also involved in corticospinal tract axon guidance independently of DCC (By similarity). Involved in dorsal root ganglion axon projection towards the spinal cord (PubMed:28483977). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (By similarity).

Cellular Location

Cell membrane; Single-pass type I membrane protein. Cell surface. Synapse, synaptosome {ECO:0000250|UniProtKB:Q761X5}. Cell projection, axon {ECO:0000250|UniProtKB:O08747}. Cell projection, dendrite {ECO:0000250|UniProtKB:O08747}. Cell projection, growth cone {ECO:0000250|UniProtKB:O08747}. Cell projection, lamellipodium {ECO:0000250|UniProtKB:O08747}. Cell projection, filopodium {ECO:0000250|UniProtKB:O08747}

Tissue Location

Mainly expressed in brain (PubMed:9782087). Expressed in temporal lobe cortical neurons and in neurons of the hippocampal pyramidal layer (PubMed:25419706). Also expressed in kidney (PubMed:9782087). Not expressed in developing or adult lung (PubMed:9782087).


Anti-UNC5C Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)

- [Cell Culture](#)

Anti-UNC5C Picoband Antibody - Images



Anti- UNC5C Picoband antibody, ABO12527, Western blottingAll lanes: Anti UNC5C (ABO12527) at 0.5ug/mlLane 1: Rat Brain Tissue Lysate at 50ugLane 2: Mouse Brain Tissue Lysate at 50ugLane 3: HELA Whole Cell Lysate at 40ugPredicted bind size: 115KDObserved bind size: 115KD

Anti-UNC5C Picoband Antibody - Background

Netrin receptor UNC5C is a protein that in humans is encoded by the UNC5C gene. This gene product belongs to the UNC-5 family of netrin receptors. Netrins are secreted proteins that direct axon extension and cell migration during neural development. They are bifunctional proteins that act as attractants for some cell types and as repellents for others, and these opposite actions are thought to be mediated by two classes of receptors. The UNC-5 family of receptors mediates the repellent response to netrin; they are transmembrane proteins containing 2 immunoglobulin (Ig)-like domains and 2 type I thrombospondin motifs in the extracellular region.