

Anti-BCAR3 Picoband Antibody
Catalog # ABO12532**Specification**

Anti-BCAR3 Picoband Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O75815
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Breast cancer anti-estrogen resistance protein 3(BCAR3) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-BCAR3 Picoband Antibody - Additional Information

Gene ID 8412

Other Names

Breast cancer anti-estrogen resistance protein 3, Novel SH2-containing protein 2, SH2 domain-containing protein 3B, BCAR3, NSP2, SH2D3B

Calculated MW

92566 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, Mouse, Rat, By Heat
Western blot, 0.1-0.5 µg/ml, Human

Tissue Specificity

Ubiquitously expressed. Found in several cancer cell lines, but not in nonmalignant breast tissue. .

Protein Name

Breast cancer anti-estrogen resistance protein 3

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg NaN₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human BCAR3 (791-825aa KGAQVNQTERYEKFNQILTALSRKLEPPPVKQAEL), different from the related mouse sequence by five amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-BCAR3 Picoband Antibody - Protein Information

Name BCAR3

Synonyms NSP2, SH2D3B

Function

Acts as an adapter protein downstream of several growth factor receptors to promote cell proliferation, migration, and redistribution of actin fibers (PubMed: [24216110](http://www.uniprot.org/citations/24216110)). Specifically involved in INS/insulin signaling pathway by mediating MAPK1/ERK2-MAPK3/ERK1 activation and DNA synthesis (PubMed: [24216110](http://www.uniprot.org/citations/24216110)). Promotes insulin-mediated membrane ruffling (By similarity). In response to vasoconstrictor peptide EDN1, involved in the activation of RAP1 downstream of PTK2B via interaction with phosphorylated BCAR1 (PubMed: [19086031](http://www.uniprot.org/citations/19086031)). Inhibits cell migration and invasion via regulation of TGFB-mediated matrix digestion, actin filament rearrangement, and inhibition of invadopodia activity (By similarity). May inhibit TGFB-SMAD signaling, via facilitating BCAR1 and SMAD2 and/or SMAD3 interaction (By similarity). Regulates EGF-induced DNA synthesis (PubMed: [18722344](http://www.uniprot.org/citations/18722344)). Required for the maintenance of ocular lens morphology and structural integrity, potentially via regulation of focal adhesion complex signaling (By similarity). Acts upstream of PTPRA to regulate the localization of BCAR1 and PTPRA to focal adhesions, via regulation of SRC-mediated phosphorylation of PTPRA (By similarity). Positively regulates integrin-induced tyrosine phosphorylation of BCAR1 (By similarity). Acts as a guanine nucleotide exchange factor (GEF) for small GTPases RALA, RAP1A and RRAS (By similarity). However, in a contrasting study, lacks GEF activity towards RAP1 (PubMed: [22081014](http://www.uniprot.org/citations/22081014)).

Cellular Location

Cytoplasm {ECO:0000250|UniProtKB:Q9QZK2}. Cell junction, focal adhesion {ECO:0000250|UniProtKB:Q9QZK2} Note=Localization to focal adhesions depends on interaction with PTPRA {ECO:0000250|UniProtKB:Q9QZK2}

Tissue Location

Ubiquitously expressed. Found in several cancer cell lines, but not in nonmalignant breast tissue


Anti-BCAR3 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Anti-BCAR3 Picoband Antibody - Images




Western blot analysis of BCAR3 expression in HEPG2 whole cell lysates (lane 1). BCAR3 at 93KD was detected using rabbit anti- BCAR3 Antigen Affinity purified polyclonal antibody (Catalog # ABO12532) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .



BCAR3 was detected in paraffin-embedded sections of mouse lymphaden tissues using rabbit anti- BCAR3 Antigen Affinity purified polyclonal antibody (Catalog # ABO12532) at 1 1/4 µg/mL. The immunohistochemical section was developed using SABC method .



BCAR3 was detected in paraffin-embedded sections of rat spleen tissues using rabbit anti- BCAR3 Antigen Affinity purified polyclonal antibody (Catalog # ABO12532) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .



BCAR3 was detected in paraffin-embedded sections of human tonsil tissues using rabbit anti-BCAR3 Antigen Affinity purified polyclonal antibody (Catalog # ABO12532) at 1 μ g/mL. The immunohistochemical section was developed using SABC method .

Anti-BCAR3 Picoband Antibody - Background

Breast cancer anti-estrogen resistance protein 3 is a protein that in humans is encoded by the BCAR3 gene. Breast tumors are initially dependent on estrogens for growth and progression and can be inhibited by anti-estrogens such as tamoxifen. However, breast cancers progress to become anti-estrogen resistant. Breast cancer anti-estrogen resistance gene 3 was identified in the search for genes involved in the development of estrogen resistance. The gene encodes a component of intracellular signal transduction that causes estrogen-independent proliferation in human breast cancer cells. The protein contains a putative src homology 2 (SH2) domain, a hall mark of cellular tyrosine kinase signaling molecules, and is partly homologous to the cell division cycle protein CDC48. Multiple transcript variants encoding different isoforms have been found for this gene.