

**Anti-CXCL5/ENA-78 Antibody**  
**Catalog # ABO12545****Specification****Anti-CXCL5/ENA-78 Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">P97885</a>
Host	Rabbit
Reactivity	Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for C-X-C motif chemokine 5(CXCL5) detection. Tested with WB, ELISA in Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-CXCL5/ENA-78 Antibody - Additional Information****Gene ID** 60665**Other Names**

C-X-C motif chemokine 5, Cytokine LIX, Small-inducible cytokine B5, Cxcl5, Scyb5

**Calculated MW**

14263 MW KDa

**Application Details**

ELISA , 0.1-0.5 µg/ml, Rat, -<br>Western blot, 0.1-0.5 µg/ml, Rat<br>

**Subcellular Localization**

Secreted.

**Protein Name**

C-X-C motif chemokine 5

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

**Immunogen**

E. coli-derived rat CXCL5 recombinant protein (Position: A38-Q130). Rat CXCL5 shares 58.7% and 76.3% amino acid (aa) sequence identity with human and mouse CXCL5, respectively.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins.

**Storage**

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Anti-CXCL5/ENA-78 Antibody - Protein Information****Name** Cxcl5**Synonyms** Scyb5**Function**

May participate in the recruitment of inflammatory cells by injured or infected tissue.


**Cellular Location**

Secreted.

**Anti-CXCL5/ENA-78 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-CXCL5/ENA-78 Antibody - Images**

Western blot analysis of CXCL5 expression in recombinant rat CXCL5 protein 0.5ng (lane 1). CXCL5 at 14KD was detected using rabbit anti- CXCL5 Antigen Affinity purified polyclonal antibody (Catalog # ABO12545) at 0.5 ??g/mL. The blot was developed using chemiluminescence (ECL)

method .

### Anti-CXCL5/ENA-78 Antibody - Background

C-X-C motif chemokine 5 or ENA-78 is a protein that in humans is encoded by the CXCL5 gene. The protein encoded by this gene, CXCL5 is a small cytokine belonging to the CXC chemokine family. It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon IFN- $\beta$ . This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2. The gene for CXCL5 is encoded on four exons and is located on human chromosome 4 amongst several other CXC chemokine genes. CXCL5 has been implicated in connective tissue remodelling. Also, CXCL5 has been described to regulate neutrophil homeostasis.