

**Anti-EGF Picoband Antibody**  
**Catalog # ABO12548****Specification****Anti-EGF Picoband Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">P01132</a>
Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Pro-epidermal growth factor(EGF) detection. Tested with WB, ELISA in Mouse.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-EGF Picoband Antibody - Additional Information****Gene ID 13645****Other Names**

Pro-epidermal growth factor, EGF, Epidermal growth factor, Egf

**Calculated MW**

133072 MW KDa

**Application Details**

ELISA , 0.1-0.5 µg/ml, Mouse, -<br>Western blot, 0.1-0.5 µg/ml, Mouse<br>

**Subcellular Localization**

Membrane; Single-pass type I membrane protein.

**Protein Name**

Pro-epidermal growth factor

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

**Immunogen**

E. coli-derived mouse EGF recombinant protein (Position: N977-R1029). Mouse EGF shares 69.8% and 77.4% amino acid (aa) sequence identity with human and rat EGF.

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins.

**Storage**

At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Anti-EGF Picoband Antibody - Protein Information****Name** Egf**Function**

EGF stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture. Magnesiotropic hormone that stimulates magnesium reabsorption in the renal distal convoluted tubule via engagement of EGFR and activation of the magnesium channel TRPM6 (By similarity).


**Cellular Location**

Membrane; Single-pass type I membrane protein.

**Anti-EGF Picoband Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-EGF Picoband Antibody - Images**

Western blot analysis of EGF expression in mouse kidney extract (lane 1). EGF at 134KD was detected using rabbit anti- EGF Antigen Affinity purified polyclonal antibody (Catalog # ABO12548) at 0.5 ??g/mL. The blot was developed using chemiluminescence (ECL) method .

**Anti-EGF Picoband Antibody - Background**

EGF is known as epidermal growth factor. This gene encodes a member of the epidermal growth factor superfamily. The encoded preproprotein is proteolytically processed to generate the 53-amino acid epidermal growth factor peptide. This protein acts a potent mitogenic factor that plays an important role in the growth, proliferation and differentiation of numerous cell types. Additionally, it acts by binding with high affinity to the cell surface receptor, epidermal growth factor receptor. Defects in this gene are the cause of hypomagnesemia type 4. Dysregulation of this gene has been associated with the growth and progression of certain cancers. Alternative splicing results in multiple transcript variants, at least one of which encodes a preproprotein that is proteolytically processed.