

Anti-Rab9 Picoband Antibody
Catalog # ABO12570**Specification**

Anti-Rab9 Picoband Antibody - Product Information

Application	WB
Primary Accession	P51151
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Ras-related protein Rab-9A(RAB9A) detection. Tested with WB in Human;Mouse;Rat.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-Rab9 Picoband Antibody - Additional Information

Gene ID 9367

Other Names

Ras-related protein Rab-9A, RAB9A, RAB9

Calculated MW

22838 MW KDa

Application Details

Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat

Subcellular Localization

Cell membrane ; Lipid-anchor ; Cytoplasmic side . Endoplasmic reticulum membrane . Golgi apparatus membrane . Late endosome. Cytoplasmic vesicle, phagosome membrane ; Lipid-anchor ; Cytoplasmic side . Cytoplasmic vesicle, phagosome. Colocalizes with OSBPL1A at the late endosome. Recruited to phagosomes containing S.aureus or M.tuberculosis.

Protein Name

Ras-related protein Rab-9A

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na₂HPO₄, 0.05mg Na₃.

Immunogen

A synthetic peptide corresponding to a sequence at the C-terminus of human Rab9 (156-193aa KDATNVAAAFEEAVRRVLATEDRSDHLIQTDTVNLHRK), different from the related mouse sequence by three amino acids, and from the related rat sequence by two amino acids.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-Rab9 Picoband Antibody - Protein Information

Name RAB9A ([HGNC:9792](#))

Synonyms RAB9

Function

The small GTPases Rab are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (By similarity). RAB9A is involved in the transport of proteins between the endosomes and the trans-Golgi network (TGN) (PubMed: [34793709](http://www.uniprot.org/citations/34793709)). Specifically uses NDE1/NDEL1 as an effector to interact with the dynein motor complex in order to control retrograde trafficking of RAB9-associated late endosomes to the TGN (PubMed: [34793709](http://www.uniprot.org/citations/34793709)). Involved in the recruitment of SGSM2 to melanosomes and is required for the proper trafficking of melanogenic enzymes TYR, TYRP1 and DCT/TYRP2 to melanosomes in melanocytes (By similarity).

Cellular Location


Cell membrane; Lipid-anchor; Cytoplasmic side. Endoplasmic reticulum membrane. Golgi apparatus membrane. Late endosome. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle membrane. Melanosome {ECO:0000250|UniProtKB:Q9R0M6}. Note=Colocalizes with OSBPL1A at the late endosome (PubMed:16176980). Recruited to phagosomes containing S.aureus or M.tuberculosis (PubMed:21255211). Mainly localizes to late endosomes and partially localizes to Golgi (PubMed:34793709) Colocalizes with NDE1 to membrane vesicles (PubMed:34793709)

Anti-Rab9 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Rab9 Picoband Antibody - Images



Western blot analysis of Rab9 expression in rat brain extract (lane 1), mouse brain extract (lane 2), HELA whole cell lysates (lane 3) and MCF-7 whole cell lysates (lane 4). Rab9 at 23KD was detected using rabbit anti- Rab9 Antigen Affinity purified polyclonal antibody (Catalog # ABO12570) at 0.5 µg/mL. The blot was developed using chemiluminescence (ECL) method .

Anti-Rab9 Picoband Antibody - Background

Ras-related protein Rab-9A, also called RAB9, is a protein that in humans is encoded by the RAB9A gene. This gene is mapped to Xp22.2. RAB9 has been localized to components of the endocytic/exocytic pathway and has been implicated in recycling of membrane receptors. It has been found that downregulation of RAB9A gene expression in HeLa cells induced severe cell vacuolation. RAB9A GTPase is directly bound by TIP47 in its active, GTP-bound conformation. Moreover, RAB9A increases the affinity of TIP47 for its cargo. What's more, this gene may involved in the transport of proteins between the endosomes and the trans Golgi network. RAB9A has been shown to interact with RABEPK and TIP47.