

Anti-SFRP4 Picoband Antibody

Catalog # ABO12574

Specification

Anti-SFRP4 Picoband Antibody - Product Information

Application WB, IHC-P, E

Primary Accession

Host
Reactivity
Clonality
Format

Polyclonal
Lyophilized

Description

Rabbit IgG polyclonal antibody for Secreted frizzled-related protein 4(SFRP4) detection. Tested with WB, IHC-P, ELISA in Human.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-SFRP4 Picoband Antibody - Additional Information

Gene ID 6424

Other Names

Secreted frizzled-related protein 4, sFRP-4, Frizzled protein, human endometrium, FrpHE, SFRP4, FRPHF

Calculated MW

39827 MW KDa

Application Details

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 μ g/ml, Human, By Heat
br> ELISA , 0.1-0.5 μ g/ml, Human, -
br> Western blot, 0.1-0.5 μ g/ml, Human
cbr>

Subcellular Localization

Secreted. Cytoplasmic in ovarian tumor cells.

Tissue Specificity

Expressed in mesenchymal cells. Highly expressed in the stroma of proliferative endometrium. Expressed in cardiomyocytes. Shows moderate to strong expression in ovarian tumors with expression increasing as the tumor stage increases. In ovarian tumors, expression levels are inversely correlated with expression of CTNNB1 (at protein level).


Protein Name

Secreted frizzled-related protein 4

Contents

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.

Immunogen





E. coli-derived human SFRP4 recombinant protein (Position: A22-K303). Human SFRP4 shares 96.8% and 96.1% amino acid (aa) sequence identity with mouse and rat SFRP4, respectively.

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins.

Storage

At -20°C for one year. After r°Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-SFRP4 Picoband Antibody - Protein Information

Name SFRP4

Synonyms FRPHE

Function

Soluble frizzled-related proteins (sFRPS) function as modulators of Wnt signaling through direct interaction with Wnts. They have a role in regulating cell growth and differentiation in specific cell types (By similarity). SFRP4 plays a role in bone morphogenesis. May also act as a regulator of adult uterine morphology and function. May also increase apoptosis during ovulation possibly through modulation of FZ1/FZ4/WNT4 signaling (By similarity). Has phosphaturic effects by specifically inhibiting sodium-dependent phosphate uptake (PubMed:12952927).

Cellular Location

Secreted. Note=Cytoplasmic in ovarian tumor cells

Tissue Location

Expressed in mesenchymal cells. Highly expressed in the stroma of proliferative endometrium. Expressed in cardiomyocytes Shows moderate to strong expression in ovarian tumors with expression increasing as the tumor stage increases. In ovarian tumors, expression levels are inversely correlated with expression of CTNNB1 (at protein level).


Anti-SFRP4 Picoband Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Anti-SFRP4 Picoband Antibody - Images





Western blot analysis of SFRP4 expression in A549 whole cell lysates (lane 1) and SW620 whole cell lysates (lane 2). SFRP4 at 49KD was detected using rabbit anti- SFRP4 Antigen Affinity purified polyclonal antibody (Catalog # ABO12574) at 0.5 ??g/mL. The blot was developed using chemiluminescence (ECL) method .



SFRP4 was detected in paraffin-embedded sections of human endometrial carcinoma tissues using rabbit anti SFRP4 Antigen Affinity purified polyclonal antibody (Catalog # ABO12574) at 1 \hat{l}_{4} g/mL. The immunohistochemical section was developed using SABC method .

Anti-SFRP4 Picoband Antibody - Background

Secreted frizzled-related protein 4 (SFRP4) is a member of the SFRP family that contains a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins. It is mapped to 7p14.1. The expression of SFRP4 in ventricular myocardium correlates with apoptosis related gene expression. And SFRP4 is a hub gene in a Type 2 Diabetes-associated gene coexpression module in human islets, and reduces glucose-induced insulin secretion through decreased \hat{l}^2 -cell exocytosis.