

**Anti-ca2 Picoband Antibody**  
**Catalog # ABO12674****Specification****Anti-ca2 Picoband Antibody - Product Information**

Application	WB, IHC-P
Primary Accession	<a href="#">P00918</a>
Host	Rabbit
Reactivity	Human, Mouse, Rat
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for Carbonic anhydrase 2(CA2) detection. Tested with WB, IHC-P in Human;Mouse;Rat.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-ca2 Picoband Antibody - Additional Information****Gene ID 760****Other Names**

Carbonic anhydrase 2, 4.2.1.1, Carbonate dehydratase II, Carbonic anhydrase C, CAC, Carbonic anhydrase II, CA-II, CA2

**Calculated MW**

29246 MW KDa

**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By Heat<br><br>Western blot, 0.1-0.5 µg/ml, Human, Mouse, Rat<br>

**Subcellular Localization**

Cytoplasm . Cell membrane . Colocalized with SLC26A6 at the surface of the cell membrane in order to form a bicarbonate transport metabolon. Displaced from the cytosolic surface of the cell membrane by PKC in phorbol myristate acetate (PMA)-induced cells.

**Protein Name**

Carbonic anhydrase 2

**Contents**

Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>.

**Immunogen**

E.coli-derived human CA2 recombinant protein (Position: S2-K260). Human CA2 shares 81.1% and 80.7% amino acid (aa) sequence identity with mouse and rat CA2, respectively.

**Purification**

Immunogen affinity purified.

### Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r° Constitution,  
at 4°C for one month. It°Can also be  
aliquotted and stored frozen at -20°C for a  
longer time.Avoid repeated freezing and  
thawing.

### Anti-ca2 Picoband Antibody - Protein Information

Name CA2

#### Function

Catalyzes the reversible hydration of carbon dioxide (PubMed:<a href="http://www.uniprot.org/citations/11327835" target="\_blank">11327835</a>, PubMed:<a href="http://www.uniprot.org/citations/11802772" target="\_blank">11802772</a>, PubMed:<a href="http://www.uniprot.org/citations/11831900" target="\_blank">11831900</a>, PubMed:<a href="http://www.uniprot.org/citations/12056894" target="\_blank">12056894</a>, PubMed:<a href="http://www.uniprot.org/citations/12171926" target="\_blank">12171926</a>, PubMed:<a href="http://www.uniprot.org/citations/1336460" target="\_blank">1336460</a>, PubMed:<a href="http://www.uniprot.org/citations/14736236" target="\_blank">14736236</a>, PubMed:<a href="http://www.uniprot.org/citations/15300855" target="\_blank">15300855</a>, PubMed:<a href="http://www.uniprot.org/citations/15453828" target="\_blank">15453828</a>, PubMed:<a href="http://www.uniprot.org/citations/15667203" target="\_blank">15667203</a>, PubMed:<a href="http://www.uniprot.org/citations/15865431" target="\_blank">15865431</a>, PubMed:<a href="http://www.uniprot.org/citations/16106378" target="\_blank">16106378</a>, PubMed:<a href="http://www.uniprot.org/citations/16214338" target="\_blank">16214338</a>, PubMed:<a href="http://www.uniprot.org/citations/16290146" target="\_blank">16290146</a>, PubMed:<a href="http://www.uniprot.org/citations/16686544" target="\_blank">16686544</a>, PubMed:<a href="http://www.uniprot.org/citations/16759856" target="\_blank">16759856</a>, PubMed:<a href="http://www.uniprot.org/citations/16807956" target="\_blank">16807956</a>, PubMed:<a href="http://www.uniprot.org/citations/17127057" target="\_blank">17127057</a>, PubMed:<a href="http://www.uniprot.org/citations/17251017" target="\_blank">17251017</a>, PubMed:<a href="http://www.uniprot.org/citations/17314045" target="\_blank">17314045</a>, PubMed:<a href="http://www.uniprot.org/citations/17330962" target="\_blank">17330962</a>, PubMed:<a href="http://www.uniprot.org/citations/17346964" target="\_blank">17346964</a>, PubMed:<a href="http://www.uniprot.org/citations/17540563" target="\_blank">17540563</a>, PubMed:<a href="http://www.uniprot.org/citations/17588751" target="\_blank">17588751</a>, PubMed:<a href="http://www.uniprot.org/citations/17705204" target="\_blank">17705204</a>, PubMed:<a href="http://www.uniprot.org/citations/18024029" target="\_blank">18024029</a>, PubMed:<a href="http://www.uniprot.org/citations/18162396" target="\_blank">18162396</a>, PubMed:<a href="http://www.uniprot.org/citations/18266323" target="\_blank">18266323</a>, PubMed:<a href="http://www.uniprot.org/citations/18374572" target="\_blank">18374572</a>, PubMed:<a href="http://www.uniprot.org/citations/18481843" target="\_blank">18481843</a>, PubMed:<a href="http://www.uniprot.org/citations/18618712" target="\_blank">18618712</a>, PubMed:<a href="http://www.uniprot.org/citations/18640037" target="\_blank">18640037</a>, PubMed:<a href="http://www.uniprot.org/citations/18942852" target="\_blank">18942852</a>, PubMed:<a href="http://www.uniprot.org/citations/1909891" target="\_blank">1909891</a>, PubMed:<a href="http://www.uniprot.org/citations/1910042" target="\_blank">1910042</a>, PubMed:<a href="http://www.uniprot.org/citations/19170619" target="\_blank">19170619</a>, PubMed:<a href="http://www.uniprot.org/citations/19186056" target="\_blank">19186056</a>, PubMed:<a href="http://www.uniprot.org/citations/19206230" target="\_blank">19206230</a>, PubMed:<a href="http://www.uniprot.org/citations/19520834" target="\_blank">19520834</a>, PubMed:<a

href="http://www.uniprot.org/citations/19778001" target="\_blank">>19778001</a>, PubMed:<a href="http://www.uniprot.org/citations/7761440" target="\_blank">>7761440</a>, PubMed:<a href="http://www.uniprot.org/citations/7901850" target="\_blank">>7901850</a>, PubMed:<a href="http://www.uniprot.org/citations/8218160" target="\_blank">>8218160</a>, PubMed:<a href="http://www.uniprot.org/citations/8262987" target="\_blank">>8262987</a>, PubMed:<a href="http://www.uniprot.org/citations/8399159" target="\_blank">>8399159</a>, PubMed:<a href="http://www.uniprot.org/citations/8451242" target="\_blank">>8451242</a>, PubMed:<a href="http://www.uniprot.org/citations/8485129" target="\_blank">>8485129</a>, PubMed:<a href="http://www.uniprot.org/citations/8639494" target="\_blank">>8639494</a>, PubMed:<a href="http://www.uniprot.org/citations/9265618" target="\_blank">>9265618</a>, PubMed:<a href="http://www.uniprot.org/citations/9398308" target="\_blank">>9398308</a>). Can also hydrate cyanamide to urea (PubMed:<a href="http://www.uniprot.org/citations/10550681" target="\_blank">>10550681</a>, PubMed:<a href="http://www.uniprot.org/citations/11015219" target="\_blank">>11015219</a>). Stimulates the chloride-bicarbonate exchange activity of SLC26A6 (PubMed:<a href="http://www.uniprot.org/citations/15990874" target="\_blank">>15990874</a>). Essential for bone resorption and osteoclast differentiation (PubMed:<a href="http://www.uniprot.org/citations/15300855" target="\_blank">>15300855</a>). Involved in the regulation of fluid secretion into the anterior chamber of the eye. Contributes to intracellular pH regulation in the duodenal upper villous epithelium during proton-coupled peptide absorption.

### Cellular Location


Cytoplasm. Cell membrane. Note=Colocalized with SLC26A6 at the surface of the cell membrane in order to form a bicarbonate transport metabolon. Displaced from the cytosolic surface of the cell membrane by PKC in phorbol myristate acetate (PMA)-induced cells

### Anti-ca2 Picoband Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### Anti-ca2 Picoband Antibody - Images



Western blot analysis of CA2 expression in rat ovary extract (lane 1), mouse liver extract (lane 2) and MCF-7 whole cell lysates (lane 3). CA2 at 29KD was detected using rabbit anti- CA2 Antigen Affinity purified polyclonal antibody (Catalog # ABO12674) at 0.5 ??g/mL. The blot was developed using chemiluminescence (ECL) method .



CA2 was detected in paraffin-embedded sections of human gastric cancer tissues using rabbit anti- CA2 Antigen Affinity purified polyclonal antibody (Catalog # ABO12674) at 1  $\frac{1}{4}$ g/mL. The immunohistochemical section was developed using SABC method .



CA2 was detected in paraffin-embedded sections of human liver cancer tissues using rabbit anti- CA2 Antigen Affinity purified polyclonal antibody (Catalog # ABO12674) at 1  $\frac{1}{4}$ g/mL. The immunohistochemical section was developed using SABC method .

#### **Anti-ca2 Picoband Antibody - Background**

CA2 is a cytosolic enzyme with the highest activity among all known CAs. The carbonic anhydrases (ACs) form a family of enzymes that catalyze the rapid interconversion of carbon dioxide and water to bicarbonate and protons (or vice versa), a reversible reaction that occurs relatively slowly in the absence of a catalyst. Mutations in the CA2 gene result in the CA II deficiency syndrome, an autosomal recessive disorder that produces osteopetrosis, renal tubular acidosis and cerebral calcification. This gene is mapped to 8q22.