

Anti-TNF alpha Antibody
Catalog # ABO12685**Specification****Anti-TNF alpha Antibody - Product Information**

Application	WB, IHC-P, E
Primary Accession	P06804
Host	Rabbit
Reactivity	Mouse
Clonality	Polyclonal
Format	Lyophilized

Description

Rabbit IgG polyclonal antibody for Tumor necrosis factor(TNF) detection. Tested with WB, IHC-P, ELISA in Mouse.

Reconstitution

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

Anti-TNF alpha Antibody - Additional Information**Gene ID 21926****Other Names**

Tumor necrosis factor, Cachectin, TNF-alpha, Tumor necrosis factor ligand superfamily member 2, TNF-a, Tumor necrosis factor, membrane form, N-terminal fragment, NTF, Intracellular domain 1, ICD1, Intracellular domain 2, ICD2, C-domain 1, C-domain 2, Tumor necrosis factor, soluble form, Tnf, Tnfa, Tnfsf2

Calculated MW
25896 MW KDa**Application Details**

Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Mouse, By Heat

ELISA , 0.1-0.5 µg/ml, Mouse, -
Western blot, 0.1-0.5 µg/ml, Mouse

Subcellular Localization

Cell membrane; Single-pass type II membrane protein.

Protein Name

tumor necrosis factor

Contents

Each vial contains 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3. Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify Carrier Free" or "No BSA" in your order note. "

Immunogen

E. coli-derived mouse TNF alpha recombinant protein(Position: L80-L235).

Purification

Immunogen affinity purified.

Cross Reactivity

No cross reactivity with other proteins

Storage

At -20°C for one year. After r° Constitution, at 4°C for one month. It° Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

Anti-TNF alpha Antibody - Protein Information**Name** Tnf**Synonyms** Tnfa, Tnfsf2**Function**

Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia. Under certain conditions it can stimulate cell proliferation and induce cell differentiation (By similarity). Induces insulin resistance in adipocytes via inhibition of insulin-induced IRS1 tyrosine phosphorylation and insulin-induced glucose uptake. Induces GKAP42 protein degradation in adipocytes which is partially responsible for TNF-induced insulin resistance (PubMed:25586176). Plays a role in angiogenesis by inducing VEGF production synergistically with IL1B and IL6 (By similarity). Promotes osteoclastogenesis and therefore mediates bone resorption (PubMed:32741026).

Cellular Location

Cell membrane; Single-pass type II membrane protein [Tumor necrosis factor, soluble form]: Secreted. [C-domain 2]: Secreted.

Anti-TNF alpha Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TNF alpha Antibody - Images



Figure. Western blot analysis of TNF alpha using anti- TNF alpha antibody (ABO12685). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V(Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane: Recombinant Mouse TNF α Protein 0.5ng. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti- TNF alpha antigen affinity purified polyclonal antibody (Catalog # ABO12685) at 0.5 μ g/mL overnight at 4 $^{\circ}$ C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for TNF alpha at approximately 17KD. The expected band size for TNF alpha is at 17KD.

Anti-TNF alpha Antibody - Background

TNF alpha(Tumor Necrosis Factor alpha) gene encodes a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor(TNF) superfamily. This cytokine is mainly secreted by macrophages. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, insulin resistance, and cancer. Knockout studies in mice also suggested the neuroprotective function of this cytokine.