

**Anti-human DDT Antibody**  
**Catalog # ABO12696****Specification****Anti-human DDT Antibody - Product Information**

Application	WB, IHC-P, ICC, E
Primary Accession	<a href="#">P30046</a>
Host	Rabbit
Reactivity	Human
Clonality	Polyclonal
Format	Lyophilized

**Description**

Rabbit IgG polyclonal antibody for D-dopachrome decarboxylase(DDT) detection. Tested with WB, IHC-P, ICC, ELISA in Human.

**Reconstitution**

Add 0.2ml of distilled water will yield a concentration of 500ug/ml.

**Anti-human DDT Antibody - Additional Information****Gene ID 1652****Other Names**

D-dopachrome decarboxylase, 4.1.1.84, D-dopachrome tautomerase, Phenylpyruvate tautomerase II, DDT

**Calculated MW**

12712 MW KDa

**Application Details**

ELISA , 0.1-0.5 µg/ml, Human, -  
Immunocytochemistry , 0.5-1 µg/ml, Human,  
-  
Immunohistochemistry(Paraffin-embedded Section), 0.5-1 µg/ml, Human, By  
Heat  
Western blot, 0.1-0.5 µg/ml, Human  
Western blot, 0.1-0.5 µg/ml, Human

**Subcellular Localization**

Cytoplasm .

**Protein Name**

D-dopachrome decarboxylase

**Contents**

Each vial contains 0.9mg NaCl, 0.2mg Na<sub>2</sub>HPO<sub>4</sub>, 0.05mg NaN<sub>3</sub>. Carrier free (No BSA) form available in stock. If you want this antibody carrier free please specify Carrier Free" or "No BSA" in your order note. "

**Immunogen**

E. coli-derived human DDT recombinant protein(Position: M1-L118).

**Purification**

Immunogen affinity purified.

**Cross Reactivity**

No cross reactivity with other proteins

Storage

At -20°C for one year. After r° Constitution, at 4°C for one month. It°Can also be aliquotted and stored frozen at -20°C for a longer time. Avoid repeated freezing and thawing.

**Anti-human DDT Antibody - Protein Information**

**Name** DDT

**Function**

Tautomerization of D-dopachrome with decarboxylation to give 5,6-dihydroxyindole (DHI).

**Cellular Location**

Cytoplasm.


**Tissue Location**

Highly expressed in the liver and at lower levels in the heart, lung and pancreas.

**Anti-human DDT Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Anti-human DDT Antibody - Images**




Figure. Western blot analysis of DDT using anti- DDT antibody (ABO12696). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions. Lane: Recombinant Human DDT Protein 0.5ng. After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti- DDT antigen affinity purified polyclonal antibody (Catalog # ABO12696) at 0.5  $\mu$ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit with Tanon 5200 system. A specific band was detected for DDT at approximately 15KD. The expected band size for DDT is at 15KD.

#### Anti-human DDT Antibody - Background

DDT, D-dopachrome tautomerization, converts D-dopachrome into 5,6-dihydroxyindole. Northern blot analysis revealed that DDT was expressed as a 0.6-kb mRNA in all tissues tested, with the strongest expression in liver. The DDT gene in human and mouse is identical in exon structure to the MIF gene. Both genes have 2 introns that are located at equivalent positions, relative to a 2-fold repeat in protein structure. The genes for DDT and MIF are closely linked on human chromosome 22 and mouse chromosome 10.